# Jørgensen's inequality for metric spaces with application to the octonions

Sarah Markham Bellerbys College 44 Cromwell Road Hove East Sussex BN3 3ER, England e-mail: smarkham@studygroup.com

John R. Parker Department of Mathematical Sciences, University of Durham, Durham DH1 3LE, England e-mail: j.r.parker@durham.ac.uk

September 10, 2004

#### Abstract

Jørgensen's inequality gives a necessary condition for the discreteness of a non-elementary group of isometries of hyperbolic 3-space. The main idea of the proof may be generalised widely but the statement is quite specialised. Here we give a scheme for restating Jørgensen's inequality for Möbius transformations of a metric space. We then show how this scheme may be applied by giving a version of Jørgensen's inequality for the octonionic hyperbolic plane.

### 1 Introduction

In [7], Jørgensen gave a famous inequality giving a necessary condition on a pair of matrices in  $SL(2, \mathbb{C})$  that generate a non-elementary discrete group. This result and its proof have been generalised to other matrix groups, see for example [13], [6], [11]. These matrix groups are all isometry groups of rank 1 symmetric spaces of non-compact type. In this paper we give a generalisation of Jørgensen's inequality to  $F_{4(-20)}$ , the isometry group of the octonionic hyperbolic plane  $\mathbf{H}^2_{\mathbb{O}}$ . In order to do so we cannot use matrix methods because of the non-associativity of the octonions.

Our approach is to consider Jørgensen's inequality not as a statement about matrices, but as a statement about Möbius transformations acting on the Riemann sphere. It is well known that a Möbius transformation can be decomposed into (orientation preserving) Euclidean isometries, dilations  $z \mapsto d^2 z$  for d > 0 and the "reflection" in the unit circle  $z \mapsto -1/z$ . (Alternatively one may use both orientation preserving and reversing Euclidean isometries, dilations and the inversion in the unit circle  $z \mapsto 1/\overline{z}$ .) The following identities are obvious.

$$|d^{2}z - d^{2}w| = d^{2}|z - w|, \ \left|\frac{1}{z}\right| = \frac{1}{|z|}, \ \left|\frac{1}{z} - \frac{1}{w}\right| = \frac{|z - w|}{|z||w|}, \ |az - z| = |a - 1||z|.$$

From the metric space viewpoint, the proof of Jørgensen's inequality only uses these four identities and well known facts about the Euclidean metric on  $\mathbb{C}$ .

In Section 2 we consider more general metric spaces X with dilations and an inversion satisfying conditions analogous to the these identities. The group generated by the isometries of X, the dilations and the inversion will be called the *Möbius group* of X, written Möb(X). We show how to extend Jørgensen's inequality to subgroups of Möb(X) with a loxodromic generator, Theorem 2.4.

In Section 3 we concentrate on the octonions. The framework of Section 2 gives us a context to generalise Jørgensen's inequality to discrete groups of octonionic hyperbolic isometries. The boundary of the octonionic hyperbolic plane  $\mathbf{H}_{\mathbb{O}}^2$  is the one point compactification of a 15 dimensional nilpotent Lie group  $\mathcal{H}^{15}$ . Moreover, by giving  $\mathcal{H}^{15}$  a particular metric, we may realise the group  $F_{4(-20)}$  as  $\text{Möb}(\mathcal{H}^{15})$ . This enables us to give a version of Jørgensen's inequality for subgroups of  $F_{4(-20)}$ with a certain type of loxodromic generator, Theorem 3.6, which is the main result of Section 3.

## 2 Jørgensen's inequality

### 2.1 Möbius transformations of a metric space

Let X be a complete metric space with metric  $\rho_0$ . Let Isom(X) be a group of isometries of X. This may be either the full isometry group or a sufficiently large subgroup, such as orientation preserving or holomorphic isometries. We will suppose that Isom(X) acts transitively on X. Let o be a distinguished point of X. (Since Isom(X) acts transitively, in fact we may take o to be any point of X.) Suppose that the stabiliser of o in Isom(X) is compact (with respect to the compact-open topology). We make some more assumptions about X that allow us to extend Isom(X) to the group of Möbius transformations on X.

Suppose that for each  $d \in \mathbb{R}_+$  there is a map  $D_d : X \longrightarrow X$  so that  $D_d o = o$  and for all  $z, w \in X$  we have

$$\rho_0(D_d z, D_d w) = d^2 \rho_0(z, w).$$
(1)

We call  $D_d$  the dilation with dilation factor  $d \in \mathbb{R}_+$ .

Let  $X \cup \{\infty\}$  be the one point compactification of X. Suppose that there is an involution R interchanging o and  $\infty$  and so that if  $z, w \in X - \{o\}$  then

$$\rho_0(Rz, o) = \frac{1}{\rho_0(z, o)},$$
(2)

$$\rho_0(Rz, Rw) = \frac{\rho_0(z, w)}{\rho_0(z, o)\rho_0(w, o)}.$$
(3)

We may think of R as reflection in the unit sphere of centre  $o \in X$ . The conditions (1), (2) and (3) generalise the first three identities given in the introduction. Let  $M\ddot{o}b(X)$  be the group generated by Isom(X),  $D_d$  and R for all  $d \in \mathbb{R}_+$ . We call  $M\ddot{o}b(X)$  the group of  $M\ddot{o}bius$  transformations of X.

**Proposition 2.1** Let X be a metric space and  $M\"{o}b(X)$  be the group generated by Isom(X),  $D_d$  and R satisfying (1), (2) and (3).

(i) Let A be any element of  $M\"{o}b(X)$  for which  $A\infty = \infty$ . Then there exists a positive number  $d_A$  so that for all  $z, w \in X$ 

$$\rho_0(Az, Aw) = d_A{}^2\rho_0(z, w).$$

(ii) Let B be any element of  $M\"{o}b(X)$  for which  $B\infty \neq \infty$ . Then there exists a positive number  $r_B$  so that for all  $z, w \in X - \{B^{-1}\infty\}$ 

$$\rho_0(Bz, Bw) = \frac{r_B^2 \rho_0(z, w)}{\rho_0(z, B^{-1} \infty) \rho_0(w, B^{-1} \infty)},$$
  
$$\rho_0(Bz, B\infty) = \frac{r_B^2}{\rho_0(z, B^{-1} \infty)}.$$

**Proof:** Any element of M"ob(X) can be written as a word in isometries of X, dilations  $D_d$  and the inversion R. We prove the result by induction on the length of these words. From (1), (2), (3) above we see that the result holds for all  $A \in \text{Isom}(X)$  (with  $d_A = 1$ ), for  $D_d$  (with  $d_{D_d} = d$ ) and for R (with  $r_R = 1$ ) since  $o = R^{-1}\infty$ .

Suppose that  $A \in \text{M\"ob}(X)$  with  $A\infty = \infty$  and suppose that A satisfies the condition (i) above. Then it is clear that  $AD_d$  and AR satisfy the conditions of (i) with  $d_{AD_d} = d_A d$  and (ii) with  $r_{AR} = d_A$  respectively.

Now suppose that  $B \in \text{M\"ob}(X)$  with  $B\infty \neq \infty$  and suppose that B satisfies the conditions of (ii) above. Consider BA,  $BD_d$ , BR for  $A \in \text{Isom}(X)$  and d > 0. Clearly  $BA\infty \neq \infty$  and  $BD_d\infty \neq \infty$ . If  $z, w \neq D_d^{-1}B^{-1}\infty$  we have

$$\rho_0(BD_d z, BD_d \infty) = \rho_0(BD_d z, B\infty)$$
$$= \frac{r_B^2}{\rho_0(D_d z, B^{-1}\infty)}$$
$$= \frac{(r_B/d)^2}{\rho_0(z, D_d^{-1}B^{-1}\infty)}$$

Similarly,

$$\rho_0(BD_d z, BD_d w) = \frac{(r_B/d)^2 \rho_0(z, w)}{\rho_0(z, D_d^{-1}B^{-1}\infty)\rho_0(w, D_d^{-1}B^{-1}\infty)}.$$

In the same way, if  $z, w \neq A^{-1}B^{-1}\infty$  we have

$$\rho_0(BAz, BA\infty) = \frac{r_B^2}{\rho_0(z, A^{-1}B^{-1}\infty)},$$
  

$$\rho_0(BAz, BAw) = \frac{r_B^2 \rho_0(z, w)}{\rho_0(z, A^{-1}B^{-1}\infty)\rho_0(w, A^{-1}B^{-1}\infty)}.$$

Suppose that  $BR\infty \neq \infty$  and  $z, w \neq RB^{-1}\infty$ . Then

$$\rho_0(BRz, BR\infty) = \rho_0(BRz, Bo) 
= \frac{r_B^2 \rho_0(Rz, o)}{\rho_0(Rz, B^{-1}\infty)\rho_0(o, B^{-1}\infty)} 
= \frac{(r_B \rho_0(RB^{-1}\infty, o))^2 \rho_0(z, o)}{\rho_0(z, o)\rho_0(z, RB^{-1}\infty)} 
= \frac{(r_B \rho_0(RB^{-1}\infty, o))^2}{\rho_0(z, RB^{-1}\infty)}.$$

Similarly

$$\rho_0(BRz, BRw) = \frac{\left(r_B\rho_0(RB^{-1}\infty, o)\right)^2 \rho_0(z, w)}{\rho_0(z, RB^{-1}\infty)\rho_0(w, RB^{-1}\infty)}$$

Finally, suppose that  $BR\infty = \infty$ , that is  $Bo = \infty$ . Then for  $z, w \neq o$  we have

$$\rho_0(BRz, BRo) = \rho_0(BRz, B\infty)$$

$$= \frac{r_B^2}{\rho_0(Rz, B^{-1}\infty)}$$

$$= \frac{r_B^2}{\rho_0(Rz, o)}$$

$$= r_B^2 \rho_0(z, o).$$

Similarly,

$$\rho_0(BRz, BRw) = r_B{}^2\rho_0(z, w)$$

Using Proposition 2.1(ii) we see that for all  $B \in M\ddot{o}b(X)$  with  $B\infty \neq \infty$  we have

$$\frac{\rho_0(Bz,z)}{\rho_0(Bz,B\infty)} = \frac{\rho_0(z,B^{-1}z)}{\rho_0(B^{-1}z,B^{-1}\infty)}.$$
(4)

Also, we may regard B as inversion in a sphere of radius  $r_B$  centred at  $B^{-1}\infty$  followed by an isometry taking  $B^{-1}\infty$  to  $B\infty$  (compare with Proposition 2.4 of [8]).

**Lemma 2.2** If Isom(X) acts transitively on X then M"ob(X) acts 2-transitively on  $X \cup \infty$ . That is, given any two pairs  $p_1$ ,  $q_1$ ;  $p_2$ ,  $q_2$  of points in  $X \cup \{\infty\}$  then there exists  $B \in G$  so that  $B(p_2) = p_1$  and  $B(q_2) = q_1$ .

**Proof:** It suffices to consider the case of  $p_2 = \infty$  and  $q_2 = o$ . Choose  $A_1$  and  $A_2$  in Isom(X) so that  $A_1o = p_1$  and  $A_2o = RA_1^{-1}q_1$ . Taking  $B = A_1RA_2$  gives

$$B\infty = A_1 R A_2 \infty = A_1 R \infty = A_1 o = p_1,$$
  
$$Bo = A_1 R A_2 o = A_1 R (R A_1^{-1} q_1) = q_1.$$

Suppose that  $A \in \text{M\"ob}(X)$  fixes  $\infty$ . From Proposition 2.1 (i), there exists  $d_A > 0$ , which we call the *dilation factor* of A, so that  $\rho_0(Az, Aw) = d_A^2 \rho_0(z, w)$ . Then A is said to be *loxodromic* if and only if  $d_A \neq 1$ . Clearly  $d_{A^{-1}} = 1/d_A$ . By the contraction mapping theorem (replacing A by  $A^{-1}$  if necessary) we see that such an A must have a unique fixed point in X. Using transitivity we may conjugate A so that this fixed point is  $o \in X$ .

An element B of  $M\ddot{o}b(X)$  is said to be *loxodromic* if it is conjugate in  $M\ddot{o}b(X)$  to a loxodromic map  $A \in M\ddot{o}b(X)$  with  $A\infty = \infty$ .

**Lemma 2.3** Suppose that  $B \in M\"{o}b(X)$  fixes distinct points  $p, q \in X \cup \{\infty\}$ . Then *B* is conjugate to  $A \in M\"{o}b(X)$  with fixed points o and  $\infty$ . Moreover, the dilation factor  $d_A$  of *A* is independent of the conjugating map.

**Proof:** The first part follows from the 2-transitivity of the action of M"ob(X) on  $X \cup \{\infty\}$ . For the second part, suppose that  $C_1BC_1^{-1} = A_1$  and  $C_2BC_2^{-1} = A_2$  where  $A_j$  fixes o and  $\infty$ . Then  $C = C_2C_1^{-1}\infty = \infty$  and so has dilation factor  $d_C$  and  $C^{-1}$  has dilation factor  $d_{C^{-1}} = 1/d_C$ . Then  $A_2 = C_2C_1^{-1}A_1C_1C_2^{-1} = CA_1C^{-1}$  and arguing as in the proof of Proposition 2.1 we see that  $A_2$  has dilation factor  $d_{A_2} = d_Cd_{A_1}d_{C^{-1}} = d_{A_1}$ .

Define the cross-ratio of quadruples of points in  $X \cup \{\infty\}$  by

$$\mathbb{X}(z_1, z_2; w_1, w_2) = \frac{\rho_0(w_1, z_1)\rho_0(w_2, z_2)}{\rho_0(w_2, z_1)\rho_0(w_1, z_2)}, \\ \mathbb{X}(z_1, z_2; \infty, w_2) = \frac{\rho_0(w_2, z_2)}{\rho_0(w_2, z_1)}.$$

Using Proposition 2.1 it is not hard to show that the cross-ratio of four points is preserved by the action of  $M\ddot{o}b(X)$ .

Let A be a loxodromic element of  $M\"{o}b(X)$  fixing  $p, q \in X \cup \{\infty\}$  and with dilation factor  $d_A$ . Suppose that  $m_A$  is a positive number so that for all points  $z \in X \cup \{\infty\} - \{p, q\}$  we have

$$\mathbb{X}(p, Az; q, z) \le d_A m_A.$$
(5)

This is a conjugation invariant statement of the following inequality in the special case when p = o and  $q = \infty$ :

$$\rho_0(z, Az) \le d_A m_A \rho_0(o, z). \tag{6}$$

This inequality generalises the fourth identity in the introduction. Observe that combining (6) with Proposition 2.1 gives

$$\rho_0(z, A^{-1}z) \le d_A^{-1}m_A\rho(z, o)$$

and so  $m_{A^{-1}} = m_A$ . We remark that such an  $m_A$  always exists. For example using Ao = o and the triangle inequality, we obtain

$$\rho_0(z, Az) \le \rho_0(z, o) + \rho_0(o, Az) = d_A(d_A + 1/d_A)\rho_0(z, o).$$

Thus one may always take  $m_A = d_A + 1/d_A \ge 2$ . However, for Jørgensen's inequality we will only be interested in loxodromic maps with  $m_A < 1$ .

#### 2.2 The main theorem

The main result of Section 2 is the following version of Jørgensen's inequality for groups of Möbius transformations on a metric space X, as defined above.

**Theorem 2.4** Let X be a complete metric space and suppose that Isom(X) acts transitively on X with compact stabilisers. Suppose that  $M\"{o}b(X)$ , the group of M\"{o}bius transformations on X, satisfies hypotheses (1), (2) and (3). Let A be a loxodromic element of  $M\"{o}b(X)$  with fixed points p and q and let  $m_A$  be a positive number satisfying (5). If  $\Gamma$  is a discrete subgroup of  $M\"{o}b(X)$  containing A, then for all  $B \in \Gamma$  so that B neither fixes nor interchanges p and q we have

$$m_A^2\Big(\mathbb{X}(Bp,q;p,Bq)+1\Big) \ge 1.$$
(7)

We now show that Theorem 2.4 generalises the case of a loxodromic generator in Jørgensen's original inequality and some of its generalisations.

#### Examples

(i) Let X be C and ρ<sub>0</sub> be the Euclidean metric. We take Isom(C) to be C ⋊ U(1), the group of holomorphic (or equivalently orientation preserving) isometries. Then C ∪ {∞} is the Riemann sphere and Möb(C) is PSL(2, C) acting by Möbius transformations. The conditions of Proposition 2.1 follow easily, as in Section 1. If λ ∈ C − {0} then set

$$Az = \frac{\lambda z}{\lambda^{-1}}, \quad Bz = \frac{az+b}{cz+d}$$

with ad - bc = 1 then  $d_A = |\lambda|$  and  $m_A = |\lambda - \lambda^{-1}|$ . The condition (7) of Theorem 2.4 becomes

$$1 \leq |\lambda - \lambda^{-1}|^2 \left( \frac{|b/d|}{|b/d - a/c|} + 1 \right) \\ = |\lambda - \lambda^{-1}|^2 (|bc| + 1) \\ = |\operatorname{tr}(A)^2 - 4| + |\operatorname{tr}[A, B] - 2|.$$

This is just the original statement of Jørgensen [7] (see also equation (1.21) of [9]).

(ii) Let X be the quaternions  $\mathbb{H}$  with the Euclidean metric  $\rho_0$ . Kellerhals gives conditions on quaternions a, b, c, d so that  $Bz = (az + b)(cz + d)^{-1}$  is a Möbius transformation (Lemma 1 of [9]). Let  $\mu$  and  $\nu$  be any purely imaginary quaternions, l > 1 be a real number and  $\alpha$ ,  $\beta$  be angles in  $[0, \pi]$  then take

$$Az = e^{l/2} \exp(\mu \alpha) z \left( e^{-l/2} \exp(-\nu \beta) \right)^{-1}, \quad Bx = (az+b)(cz+d)^{-1}$$

Clearly  $d_A = e^{l/2}$  and Kellerhals shows (in the proofs of Propositions 3 and 4 of [9]) that  $m_A = (2\cosh(l) - 2\cos(\alpha + \beta))^{1/2}$ ,  $\mathbb{X}(B\infty, o; \infty, Bo) = |bc|$ . The condition (7) becomes

$$(2\cosh(l) - \cos(\alpha + \beta))(|bc| + 1) \ge 1$$

which is equation (1.23) of [9]. Thus Theorem 2.4 is a restatement of [9] Proposition 3.

(iii) Let  $X = \mathbb{R}^n$  with the Euclidean metric  $\rho_0$  and let  $\operatorname{Isom}(\mathbb{R}^n) = \mathbb{R}^n \rtimes \operatorname{SO}(n)$ . Then we may represent elements of  $\operatorname{M\"ob}(\mathbb{R}^n)$  as  $Bz = (az+b)(cz+d)^{-1}$  where B is in  $\operatorname{SL}(\Gamma_n)$ , the group of  $2 \times 2$  matrices with elements in the Clifford group  $\Gamma_n$  satisfying Ahlfors' conditions, Definition 2.1 of [1] (see also Waterman [13]). For  $\lambda \in \Gamma_n - \{0\}$  and a, b, c, d satisfying Ahlfors' conditions, take

$$Az = \lambda z \lambda^*, \quad Bz = (az+b)(cz+d)^{-1}.$$

Then, as above  $d_A = |\lambda|$  and  $m_A = \sup |\lambda - x\lambda^{*-1}x^{-1}|$  where the supremum is taken over all  $x \in \mathbb{R}^n - \{0\}$  (compare Lemma 6.20 of [5]). Waterman (Theorem 9 of [13]) defines this quantity to be  $|\lambda - \tilde{\lambda}/|\lambda|^2|$ . Moreover, (2) and (3) are just restatements of Theorem 7(iv) and Lemma 6(i) of [13], respectively. Also, using Ahlfors' conditions, it is easy to show that

$$\mathbb{X}(B\infty, o; \infty, Bo) = \frac{|bd^{-1}|}{|ac^{-1} - bd^{-1}|} = \frac{|bd^{-1}|}{|(ad^* - bc^*)c^{*-1}d^{-1}|} = |bc|$$

Then the condition (7) of Theorem 2.4 is

$$\left|\lambda - \widetilde{\lambda}/|\lambda|^2\right|^2 (|bc|+1) \ge 1$$

which is Theorem 9(i) of [13] (see also Theorem 7.4 of [5]).

(iv) Let X be the Heisenberg group  $\mathcal{N}$  and  $\rho_0$  be the Cygan metric. We take  $\operatorname{Isom}(\mathcal{N}) = \mathcal{N} \rtimes \mathrm{U}(1)$  (which has index 2 in the full group of Heisenberg isometries). Then elements of  $\operatorname{M\"ob}(\mathcal{N})$  lie in  $\operatorname{PU}(2,1)$ . Let

$$A(\zeta, v) = \left(\lambda\zeta, |\lambda|^2 v\right)$$

for  $\lambda \in \mathbb{C} - \{0\}$ . Then we have  $d_A = |\lambda|^{1/2}$  and  $m_A = (|\lambda - 1| + |\lambda^{-1} - 1|)^{1/2}$ , see Lemma 2.1 of [12]. Also, (2) and (3) are Proposition 2.4 of [8]. Then the condition (7) of Theorem 2.4 becomes

$$\left(|\lambda - 1| + |\lambda^{-1} - 1|\right) \left(\mathbb{X}(B\infty, o; \infty, Bo) + 1\right) \ge 1$$

which is the first statement of Theorem 4.1 of [6], noting that the cross-ratio  $\mathbb{X}(z_1, z_2; w_1, w_2)$  defined above is the square root of  $|[z_1, z_2; w_1, w_2]|$  defined in [6] (see also [4] where the cross-ratio is defined in terms of the Cygan metric).

### 2.3 Proof of the main theorem

Using Lemma 2.3, we suppose that A is a loxodromic map fixing o and  $\infty$ . Suppose that B is any element of  $M\ddot{o}b(X)$  neither fixing nor interchanging o and  $\infty$ . Define a sequence  $\{B_n\}$  of elements of G as follows:

$$B_0 = B, \quad B_{n+1} = B_n A B_n^{-1}$$

For simplicity let  $r_n$  denote  $r_{B_n}$ , the constant from Proposition 2.1 (ii).

Let  $p_n = B_n o$  and  $q_n = B_n \infty$  denote the fixed points of  $B_{n+1}$ . Assume for the moment that  $p_n, q_n \neq o, \infty$  for all n. Define

$$\mathbb{X}_n = \mathbb{X}(q_n, o; \infty, p_n) = \frac{\rho_0(o, p_n)}{\rho_0(p_n, q_n)}.$$

Our goal is to show that if  $m_A^2(1 + X_0) < 1$  then there is a sequence of distinct elements in  $\langle A, B \rangle$  tending to the identity.

First, we relate  $X_{n+1}$  to  $X_n$  and use this to show that  $X_n$  tends to zero as n tends to infinity.

Lemma 2.5 With the above notation

$$\rho_0(o, p_{n+1}) \leq m_A d_A \rho_0(p_{n+1}, q_n) \frac{\rho_0(o, p_n)}{\rho_0(p_n, q_n)}, \\
\frac{1}{\rho_0(p_{n+1}, q_{n+1})} \leq \frac{m_A}{d_A \rho_0(p_{n+1}, q_n)} \frac{\rho_0(o, q_n)}{\rho_0(p_n, q_n)}.$$

-

**Proof:** Using Proposition 2.1 and (6) we have

$$\rho_{0}(o, p_{n+1}) = \rho_{0}(o, B_{n}AB_{n}^{-1}o) \\
= \frac{r_{n}^{2}\rho_{0}(B_{n}^{-1}o, AB_{n}^{-1}o)}{\rho_{0}(AB_{n}^{-1}o, B_{n}^{-1}\infty)\rho_{0}(B_{n}^{-1}o, B_{n}^{-1}\infty)} \\
\leq \frac{d_{A}m_{A}r_{n}^{2}\rho_{0}(o, B_{n}^{-1}o)}{\rho_{0}(AB_{n}^{-1}o, B_{n}^{-1}\infty)\rho_{0}(B_{n}^{-1}o, B_{n}^{-1}\infty)} \\
= \frac{d_{A}m_{A}\rho_{0}(B_{n}AB_{n}^{-1}o, B_{n}\infty)\rho_{0}(o, B_{n}o)}{\rho_{0}(B_{n}o, B_{n}\infty)} \\
= m_{A}d_{A}\rho_{0}(p_{n+1}, q_{n})\frac{\rho_{0}(o, p_{n})}{\rho_{0}(p_{n}, q_{n})}.$$

We have used (4) on the penultimate line. Similarly, we have

$$\frac{1}{\rho_0(p_{n+1}, q_{n+1})} = \frac{1}{\rho_0(B_n A B_n^{-1} o, B_n A B_n^{-1} \infty)} \\
= \frac{\rho_0(A B_n^{-1} o, B_n^{-1} \infty) \rho_0(A B_n^{-1} \infty, B_n^{-1} \infty)}{r_n^2 \rho_0(A B_n^{-1} o, A B_n^{-1} \infty)} \\
\leq \frac{d_A m_A \rho_0(A B_n^{-1} o, B_n^{-1} \infty) \rho_0(o, B_n^{-1} \infty)}{d_A^2 r_n^2 \rho_0(B_n^{-1} o, B_n^{-1} \infty)} \\
= \frac{m_A \rho_0(o, B_n \infty)}{d_A \rho_0(B_n A B_n^{-1} o, B_n \infty) \rho_0(B_n o, B_n \infty)} \\
= \frac{m_A}{d_A \rho_0(p_{n+1}, q_n)} \frac{\rho_0(o, q_n)}{\rho_0(p_n, q_n)}.$$

Combining these two estimates and using the triangle inequality we have

`

$$\begin{aligned} \mathbb{X}_{n+1} &= \frac{\rho_0(o, p_{n+1})}{\rho_0(p_{n+1}, q_{n+1})} \\ &\leq m_A^2 \frac{\rho_0(o, p_n)\rho_0(o, q_n)}{\rho_0(p_n, q_n)^2} \\ &\leq m_A^2 \frac{\rho_0(o, p_n) \left(\rho_0(o, p_n) + \rho_0(p_n, q_n)\right)}{\rho_0(p_n, q_n)^2} \\ &= m_A^2 \mathbb{X}_n(1 + \mathbb{X}_n). \end{aligned}$$

Thus, if  $m_A^2(1 + \mathbb{X}_0) < 1$  we have

$$\mathbb{X}_n \le \left(m_A^2(1+\mathbb{X}_0)\right)^n \mathbb{X}_0$$

and so  $X_n$  tends to zero exponentially as n tends to infinity.

The second step in the proof is to estimate  $\rho_0(o, p_{n+1})$  and  $\rho_0(p_{n+1}, q_{n+1})$  directly. Using Lemma 2.5 we see that we must estimate  $\rho_0(p_{n+1}, q_n)$ . Since  $X_n$  tends to zero as n tends to infinity, we may take n large enough that  $X_n$  is as small as we please.

**Lemma 2.6** Suppose that n is sufficiently large that  $d_A m_A X_n < 1$ . Then

$$\frac{\rho_0(o, q_n)}{1 + d_A m_A \mathbb{X}_n} \le \rho_0(p_{n+1}, q_n) \le \frac{\rho_0(o, q_n)}{1 - d_A m_A \mathbb{X}_n}.$$

**Proof:** We prove the lower bound. The upper bound is similar. Using the triangle inequality and Lemma 2.5 we have

$$\rho_0(o, q_n) \leq \rho_0(p_{n+1}, q_n) + \rho_0(p_{n+1}, o) \\
\leq \rho_0(p_{n+1}, q_n) (1 + d_A m_A \mathbb{X}_n).$$

Combining the inequalities from Lemmas 2.5 and 2.6, we obtain

$$\rho_{0}(o, p_{n+1}) \leq \frac{d_{A}m_{A}}{1 - d_{A}m_{A}\mathbb{X}_{n}} \frac{\rho_{0}(o, q_{n})}{\rho_{0}(p_{n}, q_{n})} \rho_{0}(o, p_{n}) \\
\leq \frac{d_{A}m_{A}(1 + \mathbb{X}_{n})}{1 - d_{A}m_{A}\mathbb{X}_{n}} \rho_{0}(o, p_{n}), \\
\frac{1}{\rho_{0}(p_{n+1}, q_{n+1})} \leq \frac{m_{A}(1 + d_{A}m_{A}\mathbb{X}_{n})}{d_{A}} \frac{1}{\rho_{0}(p_{n}, q_{n})}.$$

Define

$$\delta_n = \frac{m_A(1 + \mathbb{X}_n)}{1 - d_A m_A \mathbb{X}_n}$$

and observe that  $m_A(1 + d_A m_A \mathbb{X}_n) < \delta_n$ . Since  $m_A < 1$  and  $\mathbb{X}_n$  tends to zero as n tends to infinity, we can find an integer N so that  $\delta_N < 1$ . Thus for all  $n \ge N$  we have  $\delta_n < \delta_N$  and so

$$\rho_0(o, p_{n+1}) \le \delta_N d_A \rho_0(o, p_n), \quad \text{and} \quad \rho_0(p_{n+1}, q_{n+1}) \ge \delta_N^{-1} d_A \rho_0(p_n, q_n).$$
(8)

The third and final step is to produce a convergent sequence of distinct elements of  $\langle A, B \rangle$ . Following Jørgensen, we define  $C_n = A^{-n}B_{2n}A^n$ . Observe that (8) implies that the  $B_n$ , and hence the  $C_n$ , are distinct. Denote the fixed points of  $C_n$  by  $p'_n = A^{-n}B_{2n-1}o$  and  $q'_n = A^{-n}B_{2n-1}\infty$ . Then

$$\rho_0(p'_n, o) = d_A^{-2n} \rho_0(p_{2n-1}, o), \quad \rho_0(p'_n, q'_n) = d_A^{-2n} \rho_0(p_{2n-1}, q_{2n-1}).$$

Hence, using (8), for all  $n \ge N$  we have

$$\rho_0(p'_{n+1}, o) = d_A^{-2n-2} \rho_0(p_{2n+1}, o) \\
\leq \delta_N d_A^{-2n-1} \rho_0(p_{2n}, o) \\
\leq \delta_N^2 d_A^{-2n} \rho_0(p_{2n-1}, o) \\
= \delta_N^2 \rho_0(p'_n, o) \\
\leq \delta_N^{2(n+1-N)} \rho_0(p'_N, o).$$

Likewise

$$\rho_0(p'_{n+1}, q'_{n+1}) \ge \delta_N^{-2} \rho_0(p'_n, q'_n) \ge \delta_N^{-2(n+1-N)} \rho_0(p'_N, q'_N).$$

In particular,

$$\begin{array}{lll}
\rho_0(q'_{n+1},o) &\geq & \rho_0(p'_{n+1},q'_{n+1}) - \rho_0(p'_{n+1},o) \\
&\geq & \delta_N^{-2}\rho_0(p'_n,q'_n) - \delta_N^2\rho_0(p'_n,o) \\
&\geq & \delta_N^{-2(n+1-N)}\rho_0(p'_N,q'_N) - \delta_N^{2(n+1-N)}\rho_0(p'_N,o).
\end{array}$$

Hence for  $n \geq N$  the fixed points of  $C_n$  tend exponentially to o and  $\infty$ . We claim that the  $C_n$  lie in a compact subset of M"ob(X). Hence (a subsequence of) the  $C_n$ tend to the identity (compare [4]). Since the  $C_n$  are distinct, we see that  $\langle A, B \rangle$  is not discrete. This will prove the main theorem in the case where  $p_n, q_n \neq o, \infty$ .

In order to verify the claim, observe that we may choose  $D_n$  lying in a compact subset of M"ob(X) so that  $D_n C_n D_n^{-1}$  fixes both o and  $\infty$ . Secondly, since  $C_n$  is conjugate to A, using Lemma 2.3 we see that the dilation factor of  $D_n C_n D_n^{-1}$  is  $d_A$ . Thus for all  $z, w \in X$  we have

$$\rho_0(D_n C_n D_n^{-1} A^{-1} z, D_n C_n D_n^{-1} A^{-1} w) = d_A^2 \rho_0(A^{-1} z, A^{-1} w) = \rho_0(z, w).$$

Hence  $D_n C_n D_n^{-1} A^{-1}$  is in Isom(X) and fixes *o*. By hypothesis the stabiliser of *o* in Isom(X) is compact. Hence  $C_n$  lies in a compact subset of M"ob(X) as claimed.

We need to treat the case where there is an  $N \ge 0$  for which either  $p_N$  or  $q_N$  is o or  $\infty$ , and so  $p_{N+1} = o$  or  $q_{N+1} = \infty$ . We will not use (7) but only the fact that B neither fixes nor interchanges o and  $\infty$ . Without loss of generality, suppose  $q_{N+1} = \infty$  and hence  $q_n = \infty$  for all  $n \ge N + 1$ .

Suppose  $p_n \neq o$  for all n. Consider the sequence  $B_n$  as defined above. Since  $B_n$  is conjugate to A and fixes  $\infty$  for  $n \geq N + 1$ , we have

$$\rho_{0}(p_{n+1}, o) = \rho_{0}(B_{n}AB_{n}^{-1}o, o) 
= d_{A}^{2}\rho_{0}(AB_{n}^{-1}o, B_{n}^{-1}o) 
\leq d_{A}^{3}m_{A}\rho_{0}(B_{n}^{-1}o, o) 
\leq d_{A}m_{A}\rho_{0}(B_{n}o, o) 
\leq \delta d_{A}^{2k}\rho_{0}(p_{n}, o)$$

for some integer k and some  $\delta < 1$ . Then  $A^{-kn}B_{n+1}A^{kn}$  fixes  $p''_n = A^{-kn}p_n$  and  $\infty$ . Thus  $\rho_0(p''_n, o) = d_A^{-2kn}\rho_0(p_n, o)$  and, arguing as above,

$$\rho_0(p_{n+1}'', o) \le \delta \rho_0(p_n'', o) \le \delta^{n+1-N} \rho_0(p_N'', o).$$

Thus (a subsequence of) the  $A^{-kn}B_{n+1}A^{kn}$  converges and are distinct elements of  $\langle A, B \rangle$ . Again,  $\langle A, B \rangle$  cannot be discrete.

Finally, suppose  $p_{N+1} = o$  and  $q_{N+1} = \infty$  for some  $N \ge 0$ . Thus  $B_{n+1}$  fixes both oand  $\infty$  for all  $n \ge N+1$ . Since A has precisely two fixed points, if  $B_{n+1} = B_n A B_n^{-1}$ fixes both o and  $\infty$  then  $B_n$  either fixes or interchanges o and  $\infty$ . Without loss of generality, suppose that N is the smallest index for which  $p_{N+1} = o$  and  $q_{N+1} = \infty$ . Since  $B_0$  neither fixes nor interchanges o and  $\infty$ , we may assume that  $N \ge 1$  Then  $B_N o = \infty$  and  $B_N \infty = o$  and we see that  $B_N$  has an orbit of size 2. Thus  $B_N^2$  fixes points that  $B_N$  does not. Since  $B_N$  is conjugate to A, this is a contradiction. This proves the theorem.

### 3 The octonionic hyperbolic plane

### 3.1 The octonions

The material in this section is well known, see for example Section 2 of Allcock [2] or Section 2 of Baez [3]. The octonions  $\mathbb{O}$  comprise the real vector space spanned by  $\mathbf{e}_j$ for  $j = 0, \ldots, 7$  where  $\mathbf{e}_0 = 1$  together with a non-associative multiplication defined on the basis vectors as follows and then extended to the whole of  $\mathbb{O}$  by linearity. First for  $j = 1, \ldots, 7$ 

$$\mathbf{e}_0\mathbf{e}_j=\mathbf{e}_j\mathbf{e}_0=\mathbf{e}_j,\quad \mathbf{e}_j^2=-1.$$

Secondly for j, k = 1, ..., 7 and  $j \neq k$ 

$$\mathbf{e}_j \mathbf{e}_k = -\mathbf{e}_k \mathbf{e}_j.$$

Finally

$$\mathbf{e}_j \mathbf{e}_k = \mathbf{e}_l$$

precisely when (j, k, l) is a cyclic permutation of one of the brackets:

$$(1, 2, 4), (1, 3, 7), (1, 5, 6), (2, 3, 5), (2, 6, 7), (3, 4, 6), (4, 5, 7).$$

A multiplication table is given on page 150 of [3]. We write an octonion z as

$$z = z_0 + \sum_{j=1}^7 z_j \mathbf{e}_j.$$

Define the conjugate  $\overline{z}$  of z to be

$$\overline{z} = z_0 - \sum_{j=1}^7 z_j \mathbf{e}_j.$$

Conjugation is an anti-automorphism, that is for all octonions z and w

$$\overline{(zw)} = \overline{w}\,\overline{z}.$$

The real part of z is  $\Re(z) = \frac{1}{2}(z+\overline{z})$  and the imaginary part of z is  $\Im(z) = \frac{1}{2}(z-\overline{z})$ . The modulus |z| of an octonion is the non-negative real number defined by

$$|z|^2 = \overline{z}z = z\overline{z} = z_0^2 + z_1^2 + \dots + z_7^2.$$

The modulus is multiplicative, that is |zw| = |z| |w|. Clearly |z| > 0 unless z = 0 (that is  $z_0 = z_1 = \cdots = z_7 = 0$ ). An octonion z is a unit if |z| = 1.

The following result is due to Artin (see page 471 of [2] or pages 149–150 of [3]).

**Proposition 3.1** For any octonions x and y the subalgebra with a unit generated by x and y is associative. In particular, any product of octonions that may be written in terms of just two octonions is associative.

The following identities are due to Ruth Moufang (see page 471 of [2]).

**Proposition 3.2** Suppose that x, y, z are octanions and  $\mu$  is an imaginary unit (that is  $|\mu| = 1$  and  $\Re(\mu) = 0$  so  $1 = |\mu|^2 = -\mu^2$ ). Then

$$z(xy)z = (zx)(yz), (9)$$

$$\Re((xy)z) = \Re(x(yz)) = \Re((yz)x), \tag{10}$$

$$(\mu x \overline{\mu})(\mu y) = \mu(xy) \tag{11}$$

$$(x\mu)(\overline{\mu}y\mu) = (xy)\mu \tag{12}$$

$$xy + yx = (x\overline{\mu})(\mu y) + (y\overline{\mu})(\mu x)$$
(13)

We write  $\Re(xyz)$  for any of the 3 expressions in (10).

### 3.2 Jordan algebras and the octonionic hyperbolic plane

In this section we construct the octonionic hyperbolic plane. This should be compared to the analogous construction of the octonionic projective plane given by Baez [3]. Our construction follows Allcock [2] but we use notation that is closer to that used for complex and quaternionic hyperbolic geometries in [6], [10], [11]. Let

$$\Psi = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

Let  $M(3, \mathbb{O})$  denote the real vector space of  $3 \times 3$  matrices with octonionic entries. Let  $X^*$  denote the conjugate transpose of a matrix X in  $M(3, \mathbb{O})$ . Define

$$J = \Big\{ X \in \mathcal{M}(3, \mathbb{O}) : \Psi X = X^* \Psi \Big\}.$$

Then J is closed under the Jordan multiplication

$$X * Y = \frac{1}{2}(XY + YX),$$

and so we call J the Jordan algebra associated to  $\Psi$ . Real numbers act on  $M(3, \mathbb{O})$  by multiplication of each entry of X. We define an equivalence relation on J by  $X \sim Y$  if and only if Y = kX for some non-zero real number k. Then  $\mathbb{P}J$  is defined to be the set of equivalence classes [X].

Following Allcock we define

$$\mathbb{O}_0^3 = \left\{ \mathbf{v} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} : x, y, z \text{ all lie in some associative subalgebra of } \mathbb{O} \right\}.$$

We define an equivalence relation on  $\mathbb{O}_0^3$  by  $\mathbf{v} \sim \mathbf{w}$  if  $\mathbf{w} = \mathbf{v}\alpha$  for some  $\alpha$  in the associative subalgebra of  $\mathbb{O}$  containing the entries x, y, z of  $\mathbf{v}$ . Let  $\mathbb{PO}_0^3$  be the set of equivalence classes  $[\mathbf{v}]$ . Define a map  $\pi_{\Psi} : \mathbb{O}_0^3 \longrightarrow J$  by

$$\pi_{\Psi}(\mathbf{v}) = \mathbf{v}\mathbf{v}^{*}\Psi = \begin{pmatrix} x\overline{z} & x\overline{y} & |x|^{2} \ y\overline{z} & |y|^{2} & y\overline{x} \ |z|^{2} & z\overline{y} & z\overline{x} \end{pmatrix}.$$

It is easy to check that if  $x, y, z, \alpha$  are all octonions in the same associative subalgebra then

$$\pi_{\Psi}\begin{pmatrix}x\alpha\\y\alpha\\z\alpha\end{pmatrix} = |\alpha|^2 \pi_{\Psi}\begin{pmatrix}x\\y\\z\end{pmatrix}.$$

Hence  $\pi_{\Psi}$  descends to a map  $\overline{\pi}_{\Psi} : \mathbb{PO}_0^3 \longrightarrow \mathbb{P}J$  defined by  $\overline{\pi}_{\Psi}[\mathbf{v}] = [\pi_{\Psi}(\mathbf{v})].$ 

We define an (indefinite) norm  $|\mathbf{v}|_{\Psi}$  on  $\mathbb{O}_0^3$  by

$$|\mathbf{v}|_{\Psi} = \mathbf{v}^* \Psi \mathbf{v} = \operatorname{tr}(\pi_{\Psi}(\mathbf{v})).$$

Observe that  $|\mathbf{v}\alpha|_{\Psi} = |\alpha|^2 |\mathbf{v}|_{\Psi}$  and so the sign of  $|\mathbf{v}|_{\Psi}$  is defined consistently within classes of  $\mathbb{PO}_0^3$ . Following Allcock, a model of the octonionic hyperbolic plane  $\mathbf{H}_{\mathbb{O}}^2$ 

is the subset of  $\mathbb{P}J$  comprising  $[\pi_{\Psi}(\mathbf{v})] = \overline{\pi}_{\Psi}[\mathbf{v}]$  for  $\mathbf{v} \in \mathbb{O}_0^3$  with  $|\mathbf{v}|_{\Psi} < 0$ . Likewise its boundary is subset of  $\mathbb{P}J$  comprising  $[\pi_{\Psi}(\mathbf{v})]$  for  $\mathbf{v} \in \mathbb{O}_0^3$  with  $|\mathbf{v}|_{\Psi} = 0$ .

Let  $\mathcal{H}^{15}$  be the 15 dimensional variety

$$\mathcal{H}^{15} = \Big\{ (x, y) : x, y \in \mathbb{O}, \ x + \overline{x} + |y|^2 = 0 \Big\}.$$

The following (associative) product gives  $\mathcal{H}^{15}$  the structure of a nilpotent Lie group

$$(x,y) * (w,z) = (x+w-\overline{y}z,y+z).$$

(These coordinates on  $\mathcal{H}^{15}$  are related to those given by Allcock, page 479 of [2], by  $(x, y) = (-\frac{1}{2}|\xi|^2 + \eta, \xi)$ .) Consider the map  $\psi : \mathcal{H}^{15} \cup \{\infty\} \longrightarrow J$  given by

$$\psi(x,y) = \pi_{\Psi} \begin{pmatrix} x \\ y \\ 1 \end{pmatrix} = \begin{pmatrix} x & x\overline{y} & |x|^2 \\ y & |y|^2 & y\overline{x} \\ 1 & \overline{y} & \overline{x} \end{pmatrix}$$
$$\psi(\infty) = \pi_{\Psi} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The map  $\mathbb{P}\psi: \mathcal{H}^{15} \cup \{\infty\} \longrightarrow \partial \mathbf{H}^2_{\mathbb{O}} \subset \mathbb{P}J$  defines an isomorphism between  $\mathcal{H}^{15} \cup \{\infty\}$ and Allcock's model of the boundary of the octonionic hyperbolic plane (compare with the penultimate sentence on page 483 of [2]). Henceforth, unless we specify otherwise, we identify  $\mathcal{H}^{15} \cup \{\infty\}$  with  $\partial \mathbf{H}^2_{\mathbb{O}}$ .

We define certain transformations from  $\mathcal{H}^{15} \cup \{\infty\}$  to itself. First  $\mathcal{H}^{15}$  acts transitively on itself by left translation. For  $(t, s) \in \mathcal{H}^{15}$  define (see page 479 of [2])

$$T_{(t,s)}(x,y) = (t,s) * (x,y) = (t+x-\overline{s}y,s+y), \quad T_{(t,s)}(\infty) = \infty.$$
(14)

For any imaginary unit octonion  $\mu$  we define (see page 476 or page 479 of [2])

$$S_{\mu}(x,y) = (\mu x \overline{\mu}, y \overline{\mu}), \quad S_{\mu}(\infty) = \infty.$$
(15)

We remark that, in general,  $S_{\mu}S_{\nu} \neq S_{\mu\nu}$ . The  $S_{\mu}$  generate the compact group  $\text{Spin}_7(\mathbb{R})$  (see page 476 of [2]). For any positive real number d define

$$D_d(x,y) = (d^4x, d^2y), \quad D_d(\infty) = \infty.$$
 (16)

Finally, if  $x \neq 0$  then define (see page 476 of [2])

$$R(x,y) = (\overline{x}/|x|^2, -y\overline{x}/|x|^2), \quad R(0,0) = \infty, \quad R(\infty) = (0,0).$$
(17)

Let G be the group generated by R,  $S_{\mu}$ ,  $T_{(t,s)}$  for all imaginary unit octonions  $\mu$ and all  $(t,s) \in \mathcal{H}^{15}$ . Allcock considers this group acting not on  $\mathcal{H}^{15} \cup \{\infty\}$  but on J (by abuse of notation we have kept Allcock's names for these transformations). From this point of view, Allcock shows (Theorem 4.4 (iv) of [2]) that G is precisely the automorphism group Aut(J) of J, which is known to be  $F_{4(-20)}$ , an exceptional connected, simply connected Lie group of 52 dimensions (see for example Theorem 4.5 of [2]).

However, in [2] Allcock does not mention the dilations  $D_d$ . In fact the  $D_d$  is in G for all d > 0. For example, one may easily check that if d is a positive real number then

$$D_d(x,y) = RT_{(-2/d^2, -2/d)}RT_{(-(d-1)^2/2, d-1)}RT_{(-2,2)}RT_{(-(1-d)^2/2d^2, (1-d)/d)}(x,y).$$

Passing between our action and Allcock's is slightly subtle. Pushing our action forward under  $\mathbb{P}\psi$  and Allcock's under  $\mathbb{P}$  gives the same action on  $\partial \mathbf{H}_{\mathbb{O}}^2 \subset \mathbb{P}J$ . The preimage of this action on J used by Allcock preserves the trace (Lemma 4.1 of [2]). On the other hand, we choose the preimage under  $\mathbb{P}$  so that  $\psi(\mathcal{H}^{15} \cup \{\infty\})$ is preserved. Thus in our notation  $\psi(R(x, y))$  differs from the Allcock's by multiplication by  $1/|x|^2$ . Likewise, Allcock's dilations would be given in our notation by  $d^{-4}\psi(D_d(x, y))$  and  $d^4\psi(D_d(\infty))$ . Thus, for Allcock  $\psi(o)$  and  $\psi(\infty)$  are eigenvectors of  $D_d$  whose eigenvalues are  $d^{-4}$  and  $d^4$  respectively. Since  $d \neq 1$  these are not fixed by  $D_d$ . Hence Allcock does not include them in the stabilisers of  $\psi(o)$  and  $\psi(\infty)$  in Theorem 4.4 (v), (vi) of [2]. However, considering the action on  $\mathbb{P}J$ , we see that  $D_d$  fixes  $[\psi(o)]$  and  $[\psi(\infty)]$ . So we may adapt Allcock's theorem to show that, in terms of the action of G on  $\mathcal{H}^{15} \cup \{\infty\}$ , the stabiliser of o and  $\infty$  is generated by the  $S_{\mu}$  and  $D_d$ , and so is isomorphic to  $\mathrm{Spin}_7(\mathbb{R}) \times \mathbb{R}_+$ . Likewise, in terms of its action on  $\mathcal{H}^{15} \cup \{\infty\}$ , the stabiliser in G of  $\infty$  is  $\mathcal{H}^{15} \rtimes (\mathrm{Spin}_7(\mathbb{R}) \times \mathbb{R}_+)$ .

### 3.3 The Cygan metric

Define a bilinear form on  $M(3, \mathbb{O})$  (see page 474 of [2]) by

$$\langle X|Y\rangle = \Re \operatorname{tr}(X*Y)$$

We claim that following map  $\rho_0 : \mathcal{H}^{15} \times \mathcal{H}^{15} \longrightarrow \mathbb{R}$  gives a metric, called the *Cygan* metric, on  $\mathcal{H}^{15}$ :

$$\rho_0\big((x,y),(w,z)\big) = \left\langle \psi(x,y) | \psi(w,z) \right\rangle^{1/4}.$$
(18)

Before we prove that  $\rho_0$  is a metric we investigate some of its other properties.

First, observe that, using the Moufang identity (10), we have

$$\rho_0((x,y),(w,z))^4 = |x|^2 + |w|^2 + |y|^2 |z|^2 + \Re(xw + wx) + \Re((x\overline{y})z + y(w\overline{z}) + (y\overline{x})\overline{z} + \overline{y}(z\overline{w})) = |x|^2 + |w|^2 + |y|^2 |z|^2 + 2\Re(xw) + 2\Re(x\overline{y}z + \overline{y}z\overline{w}) = |x + \overline{w}|^2 + |\overline{z}y|^2 + 2\Re((x + \overline{w})\overline{y}z) = |x + \overline{w} + \overline{z}y|^2.$$

We now show that  $T_{(t,s)}$  and  $S_{\mu}$  preserve  $\rho_0$  and that  $\rho_0$  scales by  $d^2$  under  $D_d$ .

**Proposition 3.3** Let  $(t,s) \in \mathcal{H}^{15}$ ,  $\mu$  be an imaginary unit and d be a positive real number. Then for all  $(x, y), (z, w) \in \mathcal{H}$  we have

$$\begin{split} \rho_0 \big( T_{(t,s)}(x,y), T_{(t,s)}(w,z) \big) &= \rho_0 \big( (x,y), (w,z) \big), \\ \rho_0 \big( S_\mu(x,y), S_\mu(w,z) \big) &= \rho_0 \big( (x,y), (w,z) \big), \\ \rho_0 \big( D_d(x,y), D_d(w,z) \big) &= d^2 \rho_0 \big( (x,y), (w,z) \big) \end{split}$$

### **Proof:** We have

$$\begin{split} \rho_0\big(T_{(t,s)}(x,y),T_{(t,s)}(w,z)\big) &= \left|t+x-\overline{s}y+\overline{t}+\overline{w}-\overline{z}s+(\overline{z}+\overline{s})(s+y)\right|^{1/2} \\ &= \left|t+\overline{t}+|s|^2+x+\overline{w}+\overline{z}y\right|^{1/2} \\ &= \rho_0\big((x,y),(w,z)\big) \end{split}$$

since  $t + \bar{t} + |s|^2 = 0$ .

Using the Moufang identity (9) and  $\overline{\mu} = -\mu$  we have

$$\rho_0 (S_\mu(x, y), S_\mu(w, z)) = |\mu x \overline{\mu} + \mu \overline{w} \,\overline{\mu} + (\mu \overline{z}) (y \overline{\mu})|^{1/2}$$
$$= |\mu x \overline{\mu} + \mu \overline{w} \,\overline{\mu} + \mu (\overline{z} y) \overline{\mu})|^{1/2}$$
$$= |\mu (x + \overline{w} + \overline{z} y) \overline{\mu}|^{1/2}$$
$$= \rho_0 ((x, y), (w, z)).$$

Finally

$$\rho_0 \big( D_d(x, y), D_d(w, z) \big) = |d^4 x + d^4 \overline{w} + d^2 \overline{z} d^2 y|^{1/2} = d^2 \rho_0 \big( (x, y), (w, z) \big).$$

We are now in a position to show that  $\rho_0$  is a metric.

**Proposition 3.4** The map  $\rho_0 : \mathcal{H}^{15} \times \mathcal{H}^{15} \longrightarrow \mathbb{R}$  given by

$$\rho_0((x,y),(w,z)) = \langle \psi(x,y) | \psi(w,z) \rangle = |x + \overline{w} + \overline{z}y|^{1/2}$$

is a metric.

**Proof:** Clearly  $\rho_0((x,y),(x,y)) = |x + \overline{x} + |y|^2|^{1/2} = 0$  and for  $(x,y) \neq (z,w)$  then  $\rho_0((x,y),(w,z)) = \rho_0((w,z),(x,y))$ . If  $\rho_0((x,y),(w,z)) = 0$  then  $x + \overline{w} + \overline{z}y = 0$ . Hence

$$|y - z|^2 = |y|^2 - \overline{z}y - \overline{y}z + |z|^2$$
  
=  $-x - \overline{x} - \overline{z}y - \overline{y}z - w - \overline{w}$   
=  $-2\Re(x + \overline{w} + \overline{z}y)$   
=  $0.$ 

Thus y = z and so  $x = -\overline{w} - \overline{z}y = -\overline{w} - |z|^2 = w$ . Hence (x, y) = (w, z) as required.

It remains to prove the triangle inequality. Using the fact that  $\mathcal{H}^{15}$  acts transitively on itself by left translation  $T_{(t,s)}$ , and that this action preserves  $\rho_0$ , it suffices to show that for all  $(x, y), (w, z) \in \mathcal{H}^{15}$  we have

$$\rho_0((x,y),(w,z)) \le \rho_0((x,y),o) + \rho_0(o,(w,z))$$

where  $o = (0, 0) \in \mathcal{H}^{15}$ . In order to see this, observe

$$\rho_0((x,y),(w,z)) = |x + \overline{w} + \overline{z}y|^{1/2}$$

$$\leq (|x| + |w| + |z| |y|)^{1/2}$$

$$= (|x| + |w| + \sqrt{4\Re(x)\Re(w)})^{1/2}$$

$$\leq (|x| + 2|x|^{1/2}|w|^{1/2} + |w|)^{1/2}$$

$$= \rho_0((x,y), o) + \rho_0(o, (w,z)).$$

We have seen in Proposition 3.3 that  $T_{(t,s)}$  and  $S_{\mu}$  are isometries of the metric  $\rho_0$ and that  $D_d$  satisfies (1) with  $d_{D_d} = d$ . We will take  $\operatorname{Isom}(\mathcal{H}^{15})$  to be the subgroup of G fixing  $\infty$  and preserving  $\rho_0$ . If  $A \in G$  fixes  $\infty$  then, in Allcock's normalisation, the eigenvalue of  $\psi(\infty)$  is  $d_A^4$  where  $d_A$  is the dilation factor of A. Thus, if A also preserves  $\rho_0$ , we see that  $\psi(\infty)$  is fixed by A. Using Theorem 4.4 (vi) of [2], we see that  $\operatorname{Isom}(\mathcal{H}^{15}) = \mathcal{H}^{15} \rtimes \operatorname{Spin}_7(\mathbb{R})$ . Thus  $\operatorname{Isom}(\mathcal{H}^{15})$  acts transitively on  $\mathcal{H}^{15}$ . Moreover, the stabiliser of o = (0, 0) in  $\operatorname{Isom}(\mathcal{H}^{15})$  is the compact group  $\operatorname{Spin}_7(\mathbb{R})$ .

Finally, we show that R satisfies (2) and (3). This will imply that G, the group generated by the  $T_{(t,s)}$ ,  $S_{\mu}$ ,  $D_d$  and R, is in fact  $\text{M\"ob}(\mathcal{H}^{15})$  in the sense of the previous section.

**Proposition 3.5** Writing  $o = (0,0) \in \mathcal{H}^{15}$ , for all  $(x,y), (z,w) \in \mathcal{H}^{15} - \{o\}$  we have

$$\rho_0(R(x,y),o) = \frac{1}{\rho_0((x,y),o)},$$
  

$$\rho_0(R(x,y),R(w,z)) = \frac{\rho_0((x,y),(w,z))}{\rho_0((x,y),o)\rho_0((w,z),o)}.$$

### **Proof:** Using (10) we have

$$\begin{split} \rho_0 \big( R(x,y), R(w,z) \big)^4 &= \rho_0 \big( (\overline{x}/|x|^2, -y\overline{x}/|x|^2), (\overline{w}/|w|^2, -z\overline{w}/|w|^2) \big)^2 \\ &= \left| \overline{x}/|x|^2 + w/|w|^2 + (-w\overline{z}/|w|^2)(-y\overline{x}/|x|^2) \right|^2 \\ &= \frac{1}{|x|^2} + \frac{1}{|w|^2} + \frac{|z|^2}{|w|^2} \cdot \frac{|y|^2}{|x|^2} + 2\Re\left(\frac{\overline{x}}{|x|^2} \cdot \frac{\overline{w}}{|w|^2}\right) \\ &+ 2\Re\left(\frac{\overline{x}}{|x|^2} \cdot \frac{x\overline{y}}{|x|^2} \cdot \frac{z\overline{w}}{|w|^2}\right) + 2\Re\left(\frac{x\overline{y}}{|x|^2} \cdot \frac{z\overline{w}}{|w|^2} \cdot \frac{w}{|w|^2}\right) \\ &= \frac{|x + \overline{w} + \overline{z}y|^2}{|x|^2|w|^2} \\ &= \frac{\rho_0((x,y),(w,z))^4}{\rho_0((x,y),0)^4\rho_0((w,z),0)^4}. \end{split}$$

A similar, but easier, argument gives

$$\rho_0(R(x,y),o) = \rho_0((\overline{x}/|x|^2, -y\overline{x}/|x|^2), o) = 1/|x|^{1/2} = \frac{1}{\rho_0((x,y),o)}.$$

#### 3.4 An octonionic Jørgensen's inequality

The main result (see Chapter 4 of Markham [11]) of this section is

**Theorem 3.6** Suppose that A is a loxodromic element of  $M\"{o}b(\mathcal{H}^{15})$  conjugate to  $D_d S_\mu$  for some d > 0 and some imaginary unit  $\mu$ . Denote the fixed points of A by p and q. Let  $\Gamma$  be a discrete subgroup of  $M\"{o}b(\mathcal{H}^{15})$  containing  $D_d S_\mu$ . Then for all  $B \in \Gamma$  neither fixing nor interchanging p and q we have

$$(|d^2\mu - 1| + |d^{-2}\mu - 1|) (\mathbb{X}(Bp,q;p,Bq) + 1) \ge 1.$$

This theorem will follow directly from Theorem 2.4. We have already verified some of the hypotheses of Theorem 2.4. We know that  $\text{Isom}(\mathcal{H}^{15}) = \mathcal{H}^{15} \rtimes \text{Spin}_7(\mathbb{R})$ acts transitively on  $\mathcal{H}^{15}$  with compact stabilisers. Also,  $\text{M\"ob}(\mathcal{H}^{15})$  is generated by  $T_{(t,s)}, S_{\mu}, D_d$  and R which satisfy the hypotheses (1), (2) and (3). It remains to find a constant  $m_A$  satisfying (5). In fact we find such an  $m_A$  satisfying the equivalent condition (6).

**Proposition 3.7** Let d be a positive real number and let  $\mu$  be an imaginary unit octonion. For all  $(x, y) \in \mathcal{H}^{15}$  we have

$$\rho_0 \big( D_d S_\mu(x, y), (x, y) \big) \le d \big( |d^2 \mu - 1| + |d^{-2} \mu - 1| \big)^{1/2} \rho_0 \big( (x, y), o \big).$$

**Proof:** Using Proposition 3.1 and  $|y|^2 = -x - \overline{x}$ , we have

$$\rho_{0}(D_{d}S_{\mu}(x,y),(x,y)) = \rho_{0}((d^{4}\mu x \overline{\mu}, d^{2}y \overline{\mu}),(x,y)) \\
= |d^{4}\mu x \overline{\mu} + \overline{x} + d^{2}\overline{y}(y \overline{\mu})|^{1/2} \\
= |d^{4}\mu x \overline{\mu} + \overline{x} - d^{2}x \overline{\mu} - d^{2}\overline{x} \overline{\mu}|^{1/2} \\
= |(d^{2}\mu - 1)d^{2}x \overline{\mu} + \overline{x}d^{2}\overline{\mu}(d^{-2}\mu - 1)|^{1/2} \\
\leq d(|d^{2}\mu - 1| + |d^{-2}\mu - 1|)^{1/2}|x|^{1/2} \\
= d(|d^{2}\mu - 1| + |d^{-2}\mu - 1|)^{1/2}\rho_{0}((x,y),o).$$

Therefore if  $A = D_d S_\mu$  we have  $d_A = d$  and  $m_A = (|d^2\mu - 1| + |d^{-2}\mu - 1|)^{1/2}$ , satisfying (6). Hence all the hypotheses of Theorem 2.4 are satisfied. This proves Theorem 3.6.

Using a similar method, it is not hard to show that  $m_{D_d} = |d^2 - d^{-2}|^{1/2}$  and so in this case we could have obtained an analogous result to Theorem 3.6. However, there is an obstacle to proving Theorem 2.4 for general loxodromic maps in Möb $(\mathcal{H}^{15})$ . This can be seen by considering  $A = D_d S_\mu S_\nu$ . We have

$$\rho_0 (D_d S_\mu S_\nu(x, y), (x, y)) = \rho_0 ((d^4 \mu (\nu x \overline{\nu}) \overline{\mu}, d^2 (y \overline{\nu}) \overline{\mu}), (x, y))$$
$$= |d^4 \mu (\nu x \overline{\nu}) \overline{\mu} + \overline{x} + d^2 \overline{y} ((y \overline{\nu}) \overline{\mu})|^{1/2}$$

In general, we cannot write  $\overline{y}((y\overline{\nu})\overline{\mu}) = |y|^2\overline{\nu}\,\overline{\mu}$  and then substitute  $-x - \overline{x} = |y|^2$ .

## References

- L.V. Ahlfors, Möbius transformations and Clifford numbers, Differential Geometry and Complex Analysis, ed I. Chavel & H.M. Farkas, 65–73, Springer 1985.
- [2] D.J. Allcock, Reflection groups on the octave hyperbolic plane, Journal of Algebra 213 (1998), 467–498.
- [3] J.C. Baez, The octonions, Bulletin of the American Math. Soc. 39 (2002), 145–205.
- [4] A. Basmajian & R. Miner, Discrete subgroups of complex hyperbolic motions, Inventiones Mathematicae 131 (1998), 85–136.
- [5] C. Cao & P.L. Waterman, Conjugacy invariants of Möbius groups, Quasiconformal Mappings and Analysis, ed P. Duren et al, 109–139, Springer 1998.
- [6] Y. Jiang, S. Kamiya & J.R. Parker, Jørgensen's inequality for complex hyperbolic space, Geometriae Dedicata 97 (2003), 55–80.

- T. Jørgensen, On discrete groups of Möbius transformations, American J. Maths. 98 (1976), 739–749.
- [8] S. Kamiya, On discrete subgroups of PU(1,2; C) with Heisenberg translations,
   J. London Math. Soc. 62 (2000), 827−842.
- [9] R. Kellerhals, Quaternions and some global properties of hyperbolic 5-manifolds, Canadiian J. Maths. 55 (2003) 1080–1099.
- [10] I. Kim & J.R. Parker, Geometry of quaternionic hyperbolic manifolds Math. Proc. Cambridge Phil. Soc. 135 (2003), 291–320.
- [11] S. Markham, Hypercomplex Hyperbolic Geometry, Ph.D. thesis, University of Durham, 2003.
- [12] J.R. Parker, On the stable basin theorem, Canadian Math. Bull. (to appear).
- [13] P.L. Waterman, Möbius transformations in several dimensions, Advances in Mathematics 101 (1993), 87–113.