

# Solitons III

## (2022-23)

Stefano Cremonesi (Michaelmas)

`stefano.cremonesi@durham.ac.uk`

Charlotte Sleight (Epiphany)

`charlotte.sleight@durham.ac.uk`

Department of Mathematical Sciences, Durham University

October 6, 2022

# Contents

<b>0</b>	<b>Introduction</b>	<b>3</b>
0.1	What is a soliton? . . . . .	3
0.2	The ball-and-box model . . . . .	10
<b>1</b>	<b>Waves, dispersion and dissipation</b>	<b>15</b>
1.1	Dispersion . . . . .	15
1.2	Example: the Gaussian wave packet . . . . .	18
1.3	Dissipation . . . . .	21
1.4	Summary . . . . .	22
<b>2</b>	<b>Travelling waves</b>	<b>23</b>
2.1	The KdV soliton . . . . .	24
2.2	The sine-Gordon kink . . . . .	27
2.3	A mechanical model for the sine-Gordon equation . . . . .	29
2.4	Travelling waves and 1d point particles . . . . .	32
<b>3</b>	<b>Topological lumps and the Bogomol'nyi bound</b>	<b>37</b>
3.1	The sine-Gordon kink as a topological lump . . . . .	37
3.2	The Bogomol'nyi bound . . . . .	39
3.3	Summary . . . . .	42
<b>4</b>	<b>Conservation laws</b>	<b>44</b>
4.1	The basic idea . . . . .	45
4.2	Example: conservation of energy for sine-Gordon . . . . .	45
4.3	Extra conservation laws for relativistic field equations . . . . .	46
4.4	Conserved quantities for the KdV equation . . . . .	52
4.5	The Gardner transform . . . . .	54
<b>5</b>	<b>The Bäcklund transform</b>	<b>59</b>
5.1	Definition . . . . .	60
5.2	A simple example . . . . .	60
5.3	Bäcklund transform for sine-Gordon . . . . .	62

5.4	First example: the sine-Gordon kink from the vacuum . . . . .	63
5.5	The theorem of permutability . . . . .	65
5.6	The two-soliton solution . . . . .	69
5.7	Asymptotics of multisoliton solutions . . . . .	70
5.8	The breather . . . . .	77
<b>6</b>	<b>The Hirota method</b>	<b>82</b>
6.1	Motivation . . . . .	82
6.2	KdV equation in bilinear form . . . . .	84
6.2.1	The quadratic form of KdV . . . . .	84
6.2.2	Hirota's bilinear operator . . . . .	85
6.3	Solutions . . . . .	87
6.3.1	Example: 1-soliton . . . . .	87
6.3.2	$N$ -soliton solutions . . . . .	88
6.4	Asymptotics of 2-soliton solutions and phase shifts . . . . .	91

# Chapter 0

## Introduction

### 0.1 What is a soliton?

To a **first approximation**, solitons are very special solutions of a special class of **non-linear** partial differential equations (PDEs), or ‘**wave equations**’. (We will provide a more technical definition shortly.)

You might know that field theories, or the partial differential equations (PDEs) that describe their equations of motion, have solutions which look like **waves**. Solitons are special solutions which are localised in space and therefore look like **particles**. That’s the reason for suffix -on, as in electron, proton or photon.

The **historical discovery** of solitons occurred in 1834, when a young Scottish civil engineer named **John Scott Russell** was conducting experiments to improve the design of canal barges at the Union Canal in Hermiston, near Edinburgh, see figure 1. Accidentally, a rope pulling a barge snapped, and here is what happened next in the words of John Scott Russell himself [John Scott Russell, 1845]:

“ I was observing the motion of a boat which was rapidly drawn along a narrow channel by a pair of horses, when the boat suddenly stopped - not so the mass of water in the channel which it had put in motion; it accumulated round the prow of the vessel in a state of violent agitation, then suddenly leaving it behind, rolled forward with great velocity, assuming the form of a large solitary elevation, a rounded, smooth and well-defined heap of water, which continued its course along the channel apparently without change of form or diminution of speed.

I followed it on horseback, and overtook it still rolling on at a rate of some eight or nine miles an hour, preserving its original figure some thirty feet long and a foot




Figure 1: John Scott Russell, portrayed at a later time, and an artist's impression of the initial condition of his observation in 1834 (with a liberal interpretation of a 'pair of horses').




Figure 2: A depiction of two experiments carried out by John Scott Russell to recreate the Wave of Translation and study its properties.

to a foot and a half in height. Its height gradually diminished, and after a chase of one or two miles I lost it in the windings of the channel. Such, in the month of August 1834, was my first chance interview with that singular and beautiful phenomenon which I have called the Wave of Translation.

”

*John Scott Russell*

As we will appreciate in the coming chapters, this solitary Wave of Translation behaves very differently from the ordinary waves which solve linear differential equations, which are a good approximation when interactions are small. Different linear waves can be added up (“superimposed”) to obtain any wave profile, but these different linear waves travel at different speeds which depend on their wavelengths. As a result, any localised wave profile which is the superposition of various linear waves will “disperse” and lose its shape over time, because it consists of several linear waves which travel at different speeds. Russell’s “**Wave of Translation**”, which is now called a “**soliton**” using a term coined by [Zabusky and Kruskal, 1965], behaved very differently, maintaining its shape unaltered over a surprisingly long time. Convinced that his observation was very important, John Scott Russell followed it up by a number of experiments in which he recreated his waves of translation and studied their properties, see figure 2. His results were published ten years later in the report [John Scott Russell, 1845], but




Figure 3: From left to right: Joseph Valentin Boussinesq, Diederik Korteweg and Gustav de Vries.

much to his chagrin the scientific community paid little attention.

It took a few decades before a mathematical equation that describes shallow water waves and captures the peculiar phenomenon observed by John Scott Russell was introduced. The equation was first written down by the French mathematician and physicist Joseph Valentin Boussinesq [Boussinesq, 1877], and was then independently rediscovered by the Dutch mathematicians Diederik Korteweg and Gustav de Vries [Korteweg and Vries, 1895], see figure 3. According to the principle that things in science are named after the last people to discover them, this equation is now known as the

• **KORTEWEG-DE VRIES (KdV) EQUATION (1895):**

$$\boxed{u_t + 6uu_x + u_{xxx} = 0} . \quad (0.1)$$

This is a short-hand for the partial differential equation

$$\frac{\partial u}{\partial t} + 6u \frac{\partial u}{\partial x} + \frac{\partial^3 u}{\partial x^3} = 0$$

for the real ‘field’  $u(x, t)$ , which represents the height of a wave (measured from the surface of water at rest) in one space dimension  $x$  at time  $t$ . This equation:

- describes long wavelength shallow water waves propagating in one space dimension;
- captures the properties observed by John Scott Russell;
- is a subtle limit of the equation describing real water waves propagating in one space dimension, in coordinates moving with the wave (see [Drazin and Johnson, 1989] for details if you are interested).




Figure 4: Plot of the initial condition  $u(x, 0) = 2 \operatorname{sech}^2 x$  for the KdV equation.

### REMARKS on the KdV equation:

1. Non-linear  $\implies$  Superposition principle fails.  
(Superposition principle:  $u_1, u_2$  solutions  $\implies a_1 u_1 + a_2 u_2$  solution for all constants  $a_1, a_2$ )
2. 1st order in  $t \implies$  Solution determined by **initial condition**  $u(x, 0)$ .
3. Looks simple, but hides a rich mathematical structure.

We'll start by investigating the time evolution of the localised initial condition plotted in figure 4,

$$u(x, 0) = \frac{2}{\cosh^2(x)}, \quad (0.2)$$

with the help of a computer. To gain some intuition, let's look at the KdV equation (0.1) piece by piece:

1. Drop the non-linear term  $6uu_x$ , to obtain the **LINEARISED KdV EQUATION**:

$$u_t + u_{xxx} = 0. \quad (0.3)$$

See an animation of the time evolution [here](#). The initial localised wave **disperses**, *i.e.* it spreads out to the left, and  $u \rightarrow 0$  as  $t \rightarrow +\infty$  for any fixed  $x$ .

2. Drop the dispersive term  $u_{xxx}$ , to obtain the **DISPERSIONLESS KdV EQUATION**:

$$u_t + 6uu_x = 0. \quad (0.4)$$

In this case non-linearity causes the wave to **pile up** and **break** after a finite time:  $|u_x| \rightarrow \infty$  as  $t \rightarrow \sqrt{3}/16 \simeq 0.108$ , which can be computed using the method of characteristics. Read this if you are interested in the calculation of the breaking time and see an animation of the time evolution here (the high frequency oscillations near the breaking point are an artifact of the numerical approximation).

### 3. Keep all terms in the **KdV EQUATION**:

$$u_t + 6uu_x + u_{xxx} = 0 .$$

The two previous effects cancel and we get a “**travelling wave**”, which keeps its form and just moves to the right, as you can see here.

Admittedly, the initial condition that we chose in (0.2) was very special. Generic solutions of KdV have a much more complicated behaviour (indeed equations (0.3)-(0.4) and their solutions are recovered in certain limits). Let us then experiment with a slightly more general class of initial conditions:

$$u(x, 0) = \frac{N(N+1)}{\cosh^2(x)} , \quad N > 0, \quad (0.5)$$

which reduces to the previous initial condition (0.2) if  $N = 1$ . Here are animations of the time evolution of the initial condition (0.5) under the KdV equation, for  $N$  equal to 0.5, 1, 1.5, 2, 2.5 and 3.<sup>1</sup>

These numerical experiments indicate that:<sup>2</sup>

- **$N$  integer:**  
the initial wave splits into  $N$  **solitons** moving to the right with no dispersion.
- **$N$  not integer:**  
the initial wave splits into  $[N]$  **solitons** moving to the right plus **dispersing waves**, where  $[N]$  denotes the integer part of  $N$ .
- Either way, the different solitons move at different speeds. It can be checked that

$$\begin{array}{ll} \text{SPEED} & \propto \text{HEIGHT} \\ \text{WIDTH} & \propto (\text{HEIGHT})^{-1/2} \end{array}$$

---

<sup>1</sup>Note: in this animation space has been compactified to a circle using periodic boundary conditions  $u(10, t) = u(-10, t)$ . This allows us to investigate what happens when two solitons collide. This will be briefly discussed below, and we will return to this specific feature in greater detail later.

<sup>2</sup>We will derive these results analytically later.



in agreement with John Scott Russell's empirical observations.<sup>3</sup>

One more feature is visible if one works with periodic spatial boundary conditions (BC), in which space is a circle, as was assumed in the previous animations: faster solitons catch up with and overtake slower solitons, with seemingly no final effect on their shapes! This is very surprising for a non-linear equation, for which the superposition principle does not hold. Note also that something funny happens during the overtaking: the height of the wave decreases, unlike for linear equations where different waves add up. This unusual behaviour was first observed in experiments by John Scott Russell, who was convinced that this was very important. It took a long time for the mathematics necessary to understand this phenomenon to develop and for the scientific community to fully come on board with John Scott Russell.<sup>4</sup>

We can summarize the previous observations in the following working definition of a soliton, that we will use in the rest of the course:

A **SOLITON** is a solution of a non-linear wave equation (or PDE) which:

1. IS LOCALISED  
(It's a "lump" of energy)
2. KEEPS ITS LOCALISED SHAPE OVER TIME  
(It moves with constant shape and velocity in isolation)
3. IS PRESERVED UNDER COLLISIONS WITH OTHER SOLITONS  
(If two or more solitons collide, they re-emerge from the collision with the same shapes and velocities.)

<sup>3</sup>Roughly, KdV solitons only move to the right because the limit of the physical wave equation that leads to the KdV equation involves switching to a reference frame which moves together with the fastest possible left-moving waves. Relative to that reference frame, all other waves move to the right.

<sup>4</sup>The modern revival of solitons was kickstarted by the numerical and analytical results of [Zabusky and Kruskal, 1965], who built on the earlier important numerical work of Fermi, Pasta, Ulam and Tsingou [Fermi et al., 1955]. (The paper of Fermi et al. was based on the first ever computer-aided numerical experiment, done on the MANIAC computer at Los Alamos [Porter et al., 2009]. Mary Tsingou's role in coding the problem was neglected for a long time and has only received the attention it deserves in recent years [Dauxois, 2008].)

It was universally expected at the time that in any non-linear physical system and for any initial conditions, interactions would spread the energy of the system evenly among all its degrees of freedom over time ('thermalisation' and 'equipartition of energy') and cause the system to explore all its available configurations ('ergodicity'). This process is what makes thermodynamics and statistical mechanics work.

Fermi et al. set out to study a system of non-linearly coupled oscillators numerically, with the aim of observing how thermalisation occurs. The system initially appeared to thermalise as expected, but to their great surprise they observed that it developed close-to-periodic (rather than ergodic) behaviour over longer time scales. A decade later, Zabusky and Kruskal showed that the system studied by Fermi et al. is approximated in a certain limit by the KdV equation, whose very special properties can explain the surprising behaviour of the system.




Figure 5: Contour plot of the energy density of two colliding KdV solitons, as a function of space and time. Lighter regions have higher energy density and correspond to the cores of the two solitons. We can see the trajectories of the two solitons and the phase shift induced by the collision: the faster soliton is advanced, while the slower soliton is retarded by the collision.

Watch this video (tip: turn down the volume) of water solitons created in a lab, which obey the previous defining properties to a very good approximation.

Solitons are not just objects of purely academic interest. They can appear in Nature under a variety of circumstances. For instance, here is a video of the Severn bore taken on the 2019 spring equinox: as the high tide coming from the Atlantic Ocean enters the funnel-shaped estuary of the Severn, water surges forming highly localised waves which travel (and can be surfed!) for several miles into the Bristol Channel.

### REMARKS:

- Property 3 does not mean that nothing happens to solitons which collide: as we will study towards the end of the term, the effect of the collision is to advance or retard the solitons by a so-called “phase shift”. As an example, in figure 5 we can see the trajectories of two colliding KdV solitons and the phase shifts resulting from their interaction.
- Only very special field theories (or equivalently, wave equations) admit solitons as defined above. They are called **integrable** and are usually defined in 1 space + 1 time dimensions. Property 3 is the key. (Some people use the term “integrable soliton” for the above definition, but we will stick with “soliton” in this course.)

Solitons have been studied in depth since the 1960s in relation to many contexts:

- **Applied Maths:** water waves, optical fibres, electronics, biological systems...
- **High Energy Physics:** particle physics, gauge theory, string theory...
- **Pure Maths:** special functions, algebraic geometry, spectral theory, group theory...

We will consider two main examples of integrable equations in this course:

$$\text{KdV : } \boxed{u_t + 6uu_x + u_{xxx} = 0} \quad (0.6)$$

$$\text{sine – Gordon : } \boxed{u_{tt} - u_{xx} = -\sin u} \quad (0.7)$$

- **THIS TERM:** we will study simple **pure solitons with no dispersion**.
- **NEXT TERM:** you will study “**inverse scattering**”, a powerful formalism that allows an analytical understanding of the time evolution of **generic initial conditions**.<sup>5</sup>

To get a better feel for solitons before we start, let’s consider a discrete model which displays solitons but no dispersion. This is an example of a “cellular automaton”, a zero-player game where the rules for time evolution are fixed and the only freedom is in the choice of initial condition, but in which surprisingly rich patterns can develop.<sup>6</sup>

## 0.2 The ball-and-box model

This term we will learn several analytic methods to generate single and multiple soliton solutions of non-linear differential equations like KdV, and study the properties of these solutions.

As we have seen, experimenting with these equations on a computer can be very useful to

---

<sup>5</sup>The inverse scattering formalism was designed for equations in which space is the real line, but it is also useful if space is a finite interval or a circle (periodic bc). *E.g.* a sinusoidal initial condition on a circle evolves into a train of solitons [Zabusky and Kruskal, 1965], see this animation. Here is a contour plot of the energy density, showing the trajectories of the various solitons, which after a while recombine into a sinusoidal wave, leading to the periodic behaviour discussed in footnote4.

<sup>6</sup>The most famous cellular automaton is perhaps John Conway’s Game of Life. Read about it here if you have never heard of it. If you search Conway’s game of life or cellular automata on YouTube you will enter a rabbit hole of cool videos, often accompanied by an electronic music soundtrack. Too bad that we won’t study those cellular automata further in this course, apart from the simple model which is the subject of next section.




Figure 6: A localised configuration of the ball and box model and its continuous analogue.

develop intuition about the properties of solitons. The trouble is that you need a big-ish computer for most of these numerical experiments.

Fortunately, it was realised around 1990 that many properties of continuous solitons can be mimicked by **much simpler discrete models**, which can be studied by drawing pictures with **pen and paper**. A nice and simple example is the **BALL AND BOX MODEL** of [Takahashi and Satsuma, 1990]. In this model, **space and time** are **discrete**. In particular:

- Continuous space is replaced by an infinite line of boxes, labelled by a position  $i \in \mathbb{Z}$
- At any instant  $t \in \mathbb{Z}$ , the configuration of the system is specified by filling a number of boxes with one ball each, as in figure 6.
- Time evolution  $t \rightarrow t + 1$  is governed by the

**BALL AND BOX RULE:**

Move the leftmost ball to the next empty box to its right. Repeat the process until all balls have been moved exactly once. When you are done, the system has been evolved forward one unit in time.

The ball and box rule plays the role of the PDE for continuous solitons, *e.g.*  $u_t = -6uu_x - u_{xxx}$  in the case of the KdV equation.

**EXAMPLES:**

- **1 ball:**



We see that the two solitons keep the **same shape** after the collision, but their order is reversed: the faster soliton has overcome the slower one. If we look carefully, we can also notice that the positions of the two solitons are **delayed/advanced** by a finite amount compared to the positions that each soliton would have had in the absence of the other soliton. We will call this spatial advance or delay a “**phase shift**”, which is positive for a soliton which is advanced and negative for a soliton which is retarded. In the previous example the length-3 soliton has a phase shift of  $+4$  and the length-2 soliton has a phase shift of  $-4$ . [Make sure that you understand how this phase shift is computed from the previous figure!] This is analogous to the phase shift visible in figure 5 in the scattering of continuous KdV solitons.


**\* EXERCISE:** Generalize the previous example to a length  $m$  soliton overtaking a length  $n$  soliton (with  $m > n$ ) and find a general rule for what happens. (Start with separation  $l \geq n$  between the two solitons, that is, there are  $l$  empty boxes between the two solitons in the initial configuration.) **[Ex 4]**

The ball and box model can be generalized by introducing balls of different colours. For instance, in the **2-COLOUR BALL AND BOX MODEL**, balls come in two colours (say **BLUE** and **RED**), and again each box can be filled by at most one ball, of either colour.<sup>7</sup> The time evolution  $t \rightarrow t + 1$  is governed by the

**2-COLOUR BALL AND BOX RULE:**

Move the leftmost **BLUE** ball to the next empty box to its right. Repeat the process until all **BLUE** balls have been moved exactly once. Then do the same for the **RED** balls. When all the **BLUE** and **RED** balls have been moved, the system has been evolved forward by one unit of time.

**EXAMPLE:**



**\* EXERCISE:** Can you classify solitons in the 2-colour ball and box model? **[Ex 5]**  
 What happens when solitons collide? **[Ex 7\*]**  
 (Starred exercises are for the bravest.)

Next, we will return to continuous wave equations and aim to make the phenomenon of **dis-**

<sup>7</sup>If you happen to be colour blind and this part of the note is not accessible, please let me know and I'll replace the two colours by different symbols.

**persion** more precise.

# Chapter 1

## Waves, dispersion and dissipation

The main reference for this chapter is §1.1 of the book [Drazin and Johnson, 1989].

### 1.1 Dispersion

Usually, localised waves **spread out** (“**disperse**”) as they travel. This prevents them from being solitons. Let’s understand this phenomenon first.

#### EXAMPLES:


##### 1. ADVECTION EQUATION (linear, 1st order):

$$\frac{1}{v}u_t + u_x = 0 \quad (1.1)$$

→ Solution

$$u(x, t) = f(x - vt) \quad \text{for any function } f,$$

*i.e.* a wave moving with velocity  $v$  (right-moving if  $v > 0$ , left-moving if  $v < 0$ ). The wave keeps a fixed profile  $f(\xi)$  and moves rigidly at velocity  $v$  (indeed  $\xi = x - vt$ ):





So in this case there is no dispersion, but nothing else happens either.


2. “THE” WAVE EQUATION or D’ALEMBERT EQUATION (linear, 2nd order):

$$\boxed{\frac{1}{v^2}u_{tt} - u_{xx} = 0} \quad (v > 0 \text{ wlog}) \quad (1.2)$$

→ Solution

$$u(x, t) = f(x - vt) + g(x + vt) \quad \text{for any functions } f, g,$$

i.e. the superposition of a right-moving and a left-moving wave with velocities  $\pm v$ :



All waves move at the **same** speed, so there is no dispersion, but there is no interaction either, so this is also not very interesting for our purposes.

3. KLEIN-GORDON EQUATION<sup>1</sup> (linear, 2nd order):

$$\boxed{\frac{1}{v^2}u_{tt} - u_{xx} + m^2u = 0}, \quad (1.3)$$

where we take  $v > 0$  wlog.

This is a more interesting equation. Let us try a complex “**plane wave**” solution<sup>2</sup>

$$\boxed{u(x, t) = e^{i(kx - \omega t)}}. \quad (1.4)$$

Substituting the plane wave (1.4) in the Klein-Gordon equation (1.3), we find:

$$\begin{aligned} -\frac{\omega^2}{v^2}e^{i(kx - \omega t)} + k^2e^{i(kx - \omega t)} + m^2e^{i(kx - \omega t)} &= 0 \\ \implies -\frac{\omega^2}{v^2} + k^2 + m^2 &= 0. \end{aligned}$$

<sup>1</sup>This is the first relativistic wave equation (with  $v$  the speed of light). It was introduced independently by Oskar Klein [Klein, 1926] and Walter Gordon [Gordon, 1926], who hoped that their equation would describe electrons. It doesn’t, but it describes massive elementary particles without spin, like the pion or the Higgs boson.

<sup>2</sup>This is called a “plane wave” because its three-dimensional analogue  $u(\vec{x}, t) = \exp[i(\vec{k} \cdot \vec{x} - \omega t)]$  has constant  $u$  along a plane  $\vec{k} \cdot \vec{x} = \text{const}$  at fixed  $t$ . Unless specified, in this course we are interested in **real fields**  $u$ . It is nevertheless convenient to use complex plane waves (1.4) and eventually take the real or imaginary part to find a real solution, rather than working with the real plane waves  $\cos(kx - \omega t)$  and  $\sin(kx - \omega t)$  from the outset.

So the plane wave (1.4) is a solution of the Klein-Gordon equation (1.3) provided that  $\omega$  satisfies

$$\boxed{\omega = \omega(k) = \pm v \sqrt{k^2 + m^2}}. \quad (1.5)$$

We will usually ignore the sign ambiguity and only consider the  $+$  sign in (1.5) and similar equations.<sup>3</sup>

### VOCABULARY:

$k$	<b>wavenumber</b>	$\lambda = \frac{2\pi}{k}$	<b>wavelength</b> (periodicity in $x$ )
$\omega$	<b>angular frequency</b>	$\tau = \frac{2\pi}{\omega}$	<b>period</b> (periodicity in $t$ )

A formula like (1.5) relating  $\omega$  to  $k$ : **dispersion relation**.

The maxima of a real plane wave, like for instance  $\text{Re } e^{i(kx - \omega(k)t)}$  or  $\text{Im } e^{i(kx - \omega(k)t)}$ , are called “**wave crests**”. By a slight abuse of terminology, we will refer to the wave crests of the real or imaginary part of a complex plane wave like (1.4) simply as the wave crests of the complex plane wave.

By rewriting the complex plane wave solution (1.4) of the Klein-Gordon equation as  $e^{ik(x - c(k)t)}$ , we see that its wave crests move at the velocity

$$c(k) = \frac{\omega(k)}{k} = v \sqrt{1 + \frac{m^2}{k^2}} \text{ sign}(k).$$

Plane waves with **different wavenumbers** move at **different velocities**, so if we try to make a lump of real Klein-Gordon field by superimposing different plane waves

$$\boxed{u(x, t) = \text{Re} \int_{-\infty}^{+\infty} dk f(k) e^{i(kx - \omega(k)t)}}, \quad (1.6)$$

it will **disperse**.

In fact, there are two different notions of velocity for a wave:

#### - **PHASE VELOCITY**

$$\boxed{c(k) = \frac{\omega(k)}{k}}, \quad (1.7)$$

which is the velocity of wave crests.

---

<sup>3</sup>We do not lose generality here, since we can obtain the plane wave solution with opposite  $\omega$  by taking the complex conjugate plane wave solution and sending  $k \rightarrow -k$ .

- **GROUP VELOCITY**

$$c_g(k) = \frac{d\omega(k)}{dk}, \quad (1.8)$$

which is the velocity of the lump of field while it disperses.

We will understand better the relevance of the group velocity in the next section.

**REMARK:**

The energy (and information) carried by a wave travels at the **group velocity**, not at the phase velocity. For a **relativistic wave equation** with **speed of light**  $v$ , no signals can be transmitted faster than the speed of light. So it should be the case that  $|c_g(k)| \leq v$  for all wavenumbers  $k$ , but there is no analogous bound on the phase velocity. For example, for the Klein-Gordon equation (1.3), we can calculate

- |Group velocity|:

$$|c_g(k)| = \left| \frac{d\omega(k)}{dk} \right| = \frac{v}{\sqrt{1 + \frac{m^2}{k^2}}} \leq v$$

consistently with the principles of relativity.

- |Phase velocity|:

$$|c(k)| = \left| \frac{\omega(k)}{k} \right| = v \sqrt{1 + \frac{m^2}{k^2}} \geq v,$$

which is faster than the speed of light  $v$  for all  $k$ , but this is not a problem.

## 1.2 Example: the Gaussian wave packet

The simplest example of a localised field configuration obtained by superposition of plane waves is the “GAUSSIAN WAVE PACKET”, which is obtained by choosing a Gaussian

$$f(k) = e^{-a^2(k-\bar{k})^2} \quad (a > 0, \bar{k} \in \mathbb{R})$$

in the general superposition (1.6). This represents a lump of field with

$$\begin{array}{ll} \text{average wavenumber} & \bar{k} \\ \text{spread of wavenumber} & \sim 1/a, \end{array}$$

see fig. 1.1.




Figure 1.1: Gaussian wavepacket in Fourier space.

Then  $u(x, t) = \text{Re } z(x, t)$  is a real solution of the Klein-Gordon equation, where

$$z(x, t) = \int_{-\infty}^{+\infty} dk e^{-a^2(k-\bar{k})^2} e^{i(kx - \omega(k)t)}, \quad (1.9)$$

provided that  $\omega(k) = v \sqrt{k^2 + m^2}$ .<sup>4</sup>

Since most of the integral (1.9) comes from the region  $k \approx \bar{k}$ , we can obtain a good approximation to (1.9) by Taylor expanding  $\omega(k)$  about  $k = \bar{k}$ . Expanding to first order in  $(k - \bar{k})$  we obtain

$$\begin{aligned} \omega(k) &= \omega(\bar{k}) + \omega'(\bar{k}) \cdot (k - \bar{k}) + \mathcal{O}((k - \bar{k})^2) \\ &= \omega(\bar{k}) + c_g(\bar{k}) \cdot (k - \bar{k}) + \mathcal{O}((k - \bar{k})^2) \\ &\approx \omega(\bar{k}) + c_g(\bar{k}) \cdot (k - \bar{k}), \end{aligned}$$

where in the second line we used (1.8) and in the third line we introduced a short-hand  $\approx$  to

---

<sup>4</sup> $z(x, t)$  is a complex solution of the Klein-Gordon equation. Since the Klein-Gordon equation is a linear equation with real coefficients, the complex conjugate  $z(x, t)^*$  is also a solution of the Klein-Gordon equation, as are  $\text{Re } z(x, t)$  and  $\text{Im } z(x, t)$ .

avoid writing  $\mathcal{O}((k - \bar{k})^2)$  every time. Substituting in (1.9), we find

$$\begin{aligned}
 z(x, t) &\approx \int_{-\infty}^{+\infty} dk e^{-a^2(k-\bar{k})^2} e^{i\{kx - [\omega(\bar{k}) + c_g(\bar{k}) \cdot (k-\bar{k})]t\}} \\
 &= e^{i[\bar{k}x - \omega(\bar{k})t]} \int_{-\infty}^{+\infty} dk e^{-a^2(k-\bar{k})^2} e^{i(k-\bar{k})[x - c_g(\bar{k})t]} \\
 &\stackrel{k \rightarrow k + \bar{k}}{=} e^{i[\bar{k}x - \omega(\bar{k})t]} \int_{-\infty}^{+\infty} dk e^{-a^2k^2 + ik[x - c_g(\bar{k})t]} \\
 &\stackrel{\text{complete the square}}{=} e^{i[\bar{k}x - \omega(\bar{k})t]} e^{-\frac{1}{4a^2}[x - c_g(\bar{k})t]^2} \int_{-\infty}^{+\infty} dk e^{-a^2\{k - \frac{i}{2a^2}[x - c_g(\bar{k})t]\}^2} \\
 &\stackrel{\text{Gaussian integral}}{=} \underbrace{e^{i[\bar{k}x - \omega(\bar{k})t]}}_{\text{CARRIER WAVE}} \cdot \underbrace{\frac{\sqrt{\pi}}{a} e^{-\frac{1}{4a^2}[x - c_g(\bar{k})t]^2}}_{\text{ENVELOPE}},
 \end{aligned}$$

where in the second line we factored out a plane wave with  $k = \bar{k}$ , in the third line we changed integration variable replacing  $k$  by  $k + \bar{k}$ , in the fourth line we completed the square  $Ak^2 + Bk = A(k + \frac{B}{2A})^2 - \frac{B^2}{4A}$ , and in the last line we used the Gaussian integral formula


$$\int_{-\infty+ic}^{+\infty+ic} e^{-Ak^2} = \sqrt{\frac{\pi}{A}},$$

which holds for all  $A > 0$  and  $c \in \mathbb{R}$ . The final result is the product of a:

1. **“CARRIER WAVE”:**

a plane wave moving at the **phase velocity**


$$c(\bar{k}) = \frac{\omega(\bar{k})}{\bar{k}}$$



2. **“ENVELOPE”:**

a localised profile (or “wave packet”) moving at the **group velocity**

$$c_g(\bar{k}) = \omega'(\bar{k}).$$



Click [here](#) to see an animation of a Gaussian wavepacket with a (Gaussian) envelope and a carrier wave moving at different velocities. In the animation the phase velocity is much larger than the group velocity.

To this order of approximation, the spatial **width** of the lump has the *parametric dependence*

$$\text{WIDTH} \sim a ,$$

meaning that the width doubles if  $a$  is doubled, and is constant in time. (Indeed, a simultaneous rescaling of  $x - c_g(\bar{k})t$  and  $a$  by the same constant  $\lambda$  leaves the envelope invariant.)

**\* EXERCISE:** Improve on the previous approximation by including the 2nd order in  $k - \bar{k}$ . Show that **[Ex 10]**

$$\text{WIDTH}^2 \sim a^2 + \frac{\omega''(\bar{k})}{4a^2} t^2$$

and that the amplitude of the wave packet also decreases as time increases.

This leads to the phenomenon of **DISPERSION**, whereby the profile of the wave packet changes as it propagates. In particular, starting from a localised wave packet, dispersion makes the wave packet spread out: the width of the initial wave packet grows and the amplitude decreases as time increases. See this animation of the time evolution of the Gaussian wave-packet up to second order in  $(k - \bar{k})$ .

### 1.3 Dissipation

So far we have considered wave equations which lead to a real dispersion relation, so  $\omega(k) \in \mathbb{R}$ . If instead  $\omega(k) \in \mathbb{C}$ , then a new phenomenon occurs: **DISSIPATION**, where the **amplitude** of the wave **decays (or grows) exponentially in time**. For a plane wave

$$u(x, t) = e^{i(kx - \omega(k)t)} = e^{i(kx - \text{Re } \omega(k) \cdot t)} e^{\text{Im } \omega(k) \cdot t} \quad (1.10)$$

and we have two cases:

- $\text{Im } \omega(k) < 0$ : **“PHYSICAL DISSIPATION”**  
The amplitude **decays** exponentially with time.
- $\text{Im } \omega(k) > 0$ : **“UNPHYSICAL DISSIPATION”**  
The amplitude **grows** exponentially with time (physically unacceptable).

#### EXAMPLES:

1.

$$\boxed{\frac{1}{v} u_t + u_x + \alpha u = 0} \quad (\alpha > 0) \quad (1.11)$$

Sub in a plane wave  $u = e^{i(kx - \omega t)}$ :

$$-i \frac{\omega}{v} + ik + \alpha = 0 \implies \omega(k) = v(k - i\alpha) ,$$

leading to a **complex** dispersion relation. The plane wave solution is therefore

$$u(x, t) = e^{ik(x-vt)} e^{-\alpha vt}$$

and the wave decays exponentially, or “**dissipates**”, to zero as  $t \rightarrow +\infty$ . This is an example of physical dissipation. ( $\alpha < 0$  would have led to unphysical dissipation.)

## 2. HEAT EQUATION:

$$\boxed{u_t - \alpha u_{xx} = 0} \quad (\alpha > 0) \quad (1.12)$$

\* **EXERCISE:** Sub in a plane wave and derive the dispersion relation  $\omega(k) = -i\alpha k^2$ .

So the plane wave solution of the heat equation is

$$u(x, t) = e^{ikx} e^{-\alpha k^2 t}$$

and the waves dissipates as time passes.

## 1.4 Summary

- **Linear** wave equation  $\longrightarrow$  (Complex) **plane wave** solutions  $u = e^{i(kx - \omega t)}$ .  
Sub in to get  $\omega = \omega(k)$  **dispersion relation**.
- Wave crests move at  $c(k) = \omega(k)/k$  **phase velocity**.  
(If  $\omega(k) \in \mathbb{C}$ , then we define the phase velocity as  $c(k) = \text{Re } \omega(k)/k$ .)
- Lumps of field move at  $c_g(k) = \omega'(k)$  **group velocity**.  
/wave packets  
(If  $\omega(k) \in \mathbb{C}$ , then we define the group velocity as  $c_g(k) = \text{Re } \omega'(k)$ .)
- **Dispersion** (real  $\omega$ , width increases and amplitude decreases) and **dissipation** (complex  $\omega$ , amplitude decreases exponentially) smooth out and destroy localised lumps of energy in linear wave (or field) equations.
- **Non-linearity** can have an opposite effect (steepening and breaking, see chapter 0).
- For **solitons** the competing effects counterbalance one another precisely, leading to stable lumps of energy, unlike for ordinary waves.

# Chapter 2

## Travelling waves

The main references for this chapter are §2.1-2.2 of [Drazin and Johnson, 1989] and §2.1 of [Dauxois and Peyrard, 2006].

A “**TRAVELLING WAVE**” is a solution of a wave equation of the form

$$u(x, t) = f(x - vt) ,$$

where  $f$  is a function of a single variable, which we will typically denote by  $\xi := x - vt$ . The **velocity**  $v$  of the travelling wave could either be:

1. **Fixed** in terms of a **parameter** appearing in the **wave equation**, as in d’Alembert’s general solution

$$u(x, t) = f(x - vt) + g(x + vt)$$

of the wave equation

$$\frac{1}{v^2} u_{tt} - u_{xx} = 0 ,$$

which is the linear superposition of two travelling waves with velocities  $\pm v$ .

2. A **free parameter of the solution**, as in the KdV soliton that we will derive shortly.

### **REMARK:**

In some cases (*e.g.* “the” wave equation or the sine-Gordon equation) there will be both a velocity parameter appearing in the equation (*e.g.* the speed of light) and a different velocity parameter appearing in the travelling wave solution (namely, the speed of the wave). To avoid confusion, from now on the velocity parameter appearing in the wave equation will be set to



1 by an appropriate choice of units, and  $v$  will be reserved for the velocity of the travelling wave. For example, we will write “the” wave equation as  $u_{tt} - u_{xx} = 0$  and d’Alembert’s general solution as  $u(x, t) = f(x - t) + g(x + t)$ , which is the superposition of two travelling waves with velocities  $v = \pm 1$ .

## 2.1 The KdV soliton

We would like to find a travelling wave solution of the KdV equation

$$u_t + 6uu_x + u_{xxx} = 0$$

with **boundary conditions** (BC’s)

$$\text{BC's : } \quad u, u_x, u_{xx} \xrightarrow{x \rightarrow \pm\infty} 0$$

for all finite values of  $t$ . Let us accept these BC’s for the time being; we will derive them later.

Substituting in the KdV equation the travelling wave ansatz  $u(x, t) = f(x - vt) \equiv f(\xi)$  where  $\xi = x - vt$ , using the chain rule to express partial derivatives wrt  $x$  and  $t$  in terms of ordinary derivatives wrt  $\xi$  as follows,

$$\frac{\partial}{\partial x} = \frac{\partial \xi}{\partial x} \frac{d}{d\xi} = \frac{d}{d\xi}, \quad \frac{\partial}{\partial t} = \frac{\partial \xi}{\partial t} \frac{d}{d\xi} = -v \frac{d}{d\xi},$$

and using primes to denote derivatives wrt  $\xi$ , we obtain an ODE which we can integrate twice:


$$\begin{aligned} & -vf' + 6ff' + f''' = 0 \\ \xRightarrow{\int d\xi} & -vf + 3f^2 + f'' = A \\ \xRightarrow{\int d\xi f'} & -\frac{v}{2}f^2 + f^3 + \frac{1}{2}(f')^2 = Af + B, \end{aligned}$$

where  $A$  and  $B$  are integration constants. The second integration used an integrating factor  $f'$ , as denoted by the short-hand  $\int d\xi f'$ .

We can determine the integration constants  $A$  and  $B$  by imposing the BC’s, which imply that  $f, f', f'' \rightarrow 0$  as  $\xi \rightarrow \pm\infty$ . Sending  $\xi \rightarrow \pm\infty$  in the second and third line above we find<sup>1</sup>

$$\begin{aligned} \text{BC's:} & \quad A = B = 0 \\ \Rightarrow & \quad (f')^2 = f^2(v - 2f) \\ \Rightarrow & \quad f' = \pm f\sqrt{v - 2f} \\ \Rightarrow & \quad \int \frac{df}{f\sqrt{v - 2f}} = \pm \xi \equiv \pm(x - vt). \end{aligned} \quad (*)$$

---

<sup>1</sup>  Always impose the boundary conditions carefully and keep in mind that they don’t always imply that the integration constants vanish. This is a major source of mistakes in homework and exams.

where we note that we need  $f \leq v/2$  to ensure that  $f, f' \in \mathbb{R}$ .

To calculate the integral obtained by separation of variables, we change integration variable

$$\begin{aligned}
 f &= \frac{v}{2} \operatorname{sech}^2 \vartheta & (**) \\
 \implies df &= -v \frac{\sinh \vartheta}{\cosh^3 \vartheta} d\vartheta, \\
 \sqrt{v-2f} &= \sqrt{v} \sqrt{1 - \frac{1}{\cosh^2 \vartheta}} = \pm \sqrt{v} \frac{\sinh \vartheta}{\cosh \vartheta} \\
 \implies \frac{df}{f \sqrt{v-2f}} &= \mp \frac{v \frac{\sinh \vartheta}{\cosh^3 \vartheta} d\vartheta}{\frac{v}{2} \frac{1}{\cosh^2 \vartheta} \sqrt{v} \frac{\sinh \vartheta}{\cosh \vartheta}} = \mp \frac{2}{\sqrt{v}} d\vartheta. & (***)
 \end{aligned}$$

Substituting (\*\*) in (\*) and keeping in mind that the sign ambiguities arising from taking square roots in the two equations are unrelated (and therefore only the relative sign ambiguity matters), we find

$$\begin{aligned}
 -\frac{2}{\sqrt{v}} \int d\vartheta &= \pm(x - vt) \\
 \implies \vartheta &= \pm \frac{\sqrt{v}}{2} (x - x_0 - vt),
 \end{aligned}$$

where  $x_0$  is an integration constant. Substituting (\*\*) we find the travelling wave solution

$$\boxed{u(x, t) = f(x - vt) = \frac{v}{2} \operatorname{sech}^2 \left[ \frac{\sqrt{v}}{2} (x - x_0 - vt) \right]} \quad (2.1)$$

where the sign ambiguity has disappeared because  $\operatorname{sech}^2$  is an even function.

The travelling wave solution (2.1) of the KdV equation is the **KdV SOLITON**. See 2.1 for a snapshot of the KdV soliton.

### REMARKS:

- For a **real non-singular** solution we need  $v \geq 0$ , which means that KdV solitons only travel to the right.<sup>2</sup>

---

<sup>2</sup>If  $v < 0$  the travelling wave solution is

$$-\frac{|v|}{2} \operatorname{sech}^2 \left[ \frac{\sqrt{|v|}}{2} (x - x_0 + |v|t) \right],$$

which moves to the left with speed  $|v|$  but **diverges** wherever  $[\dots] = (n + \frac{1}{2})\pi$  with  $n \in \mathbb{Z}$ . We are always after real bounded solutions, so we discard this singular (or divergent) solution.




Figure 2.1: Snapshot of the KdV soliton.

- **PROPERTIES** of the KdV soliton:

VELOCITY	$v$
HEIGHT	$v/2$
WIDTH	$\sim \frac{1}{\sqrt{v}}$
CENTRE	$x_0 + vt$

### Clarification:

What do I mean by WIDTH  $\sim 1/\sqrt{v}$ ? A **possible** definition of the width of the soliton is as the distance between the two points where the value of  $u$  is reduced by a factor of  $e$  from its maximum, that is WIDTH  $= |x_+ - x_-| \equiv 2\Delta x$  where  $u(x_{\pm}) = v/(2e)$ . For  $\sqrt{v}\Delta x \gg 1$ , we can approximate  $\text{sech}^2\left(\frac{\sqrt{v}}{2}\Delta x\right) \approx 4e^{-\sqrt{v}\Delta x}$ , therefore this definition of width would give

$$\text{WIDTH} = 2\Delta x \approx \frac{2}{\sqrt{v}}(1 + \log 4) \approx \frac{4.77}{\sqrt{v}}.$$

(Without the approximation one finds  $4.34.../\sqrt{v}$ .) However the above definition of width was somewhat arbitrary: for instance we could have looked at points where the value  $u$  is reduced by a factor of 2, or 3, or else, from its maximum. Given a precise definition of width, one can determine the precise coefficient of  $1/\sqrt{v}$  above, but fixating on a precise definition would be somewhat absurd given the arbitrariness in the definition. It is better to say that **“the width is of the order of”** (or equivalently **“goes like”**)  $1/\sqrt{v}$ . This is independent of the precise definition of width and captures the essential point that the spatial coordinate  $x$  appears multiplied by  $\sqrt{v}$  in the KdV soliton solution (2.1). We use  $\sim$  to denote this **parametric dependence**. This is not to be confused with  $\approx$ , which means “is approximately equal to”.

A final comment: if the BC's are changed to allow  $A, B \neq 0$  (e.g. if we impose periodic boundary conditions, which is equivalent to solving the KdV equation on a circle), then the ODE for the travelling wave solution can still be integrated exactly using elliptic functions. See §2.4, 2.5 of [Drazin and Johnson, 1989] if you are interested.

## 2.2 The sine-Gordon kink

Let us seek a travelling wave solution the **sine-Gordon** equation

$$u_{xx} - u_{tt} = \sin u ,$$

where  $u$  is an angular variable  $u$  defined modulo  $2\pi$ , subject to the boundary conditions

$$\text{BC's :} \quad u \bmod 2\pi, \quad u_x \xrightarrow{x \rightarrow \pm\infty} 0$$

for every finite  $t$ . (More about these BC's later.)

Substituting the travelling wave ansatz  $u(x, t) = f(x - vt) \equiv f(\xi)$  in the sine-Gordon equation, we find

$$\begin{aligned} (1 - v^2)f'' &= \sin f \\ \iff f'' &= \gamma^2 \sin f, \quad \text{where } \gamma := \frac{1}{\sqrt{1 - v^2}} \\ \xRightarrow{\int d\xi f'} \frac{1}{2}(f')^2 &= A - \gamma^2 \cos f \\ \text{BC's:} \quad A &= \gamma^2 \\ \implies f' &= \pm \sqrt{2}\gamma \sqrt{1 - \cos f} = \pm 2\gamma \sin \frac{f}{2} \\ \implies \int \frac{df}{2 \sin \frac{f}{2}} &= \pm \gamma(x - x_0 - vt) \\ \implies \log \tan \frac{f}{4} &= \pm \gamma(x - x_0 - vt) \end{aligned}$$

where  $x_0$  is an undetermined integration constant.

We find therefore the following travelling wave solution of the sine-Gordon equation

$$\boxed{u(x, t) = f(x - vt) = 4 \arctan \left( e^{\pm \gamma(x - x_0 - vt)} \right)}, \quad (2.2)$$

which goes by the name of “**KINK**” (+ sign) or “**ANTI-KINK**” (− sign).

Note that the BC required that as  $\xi \rightarrow \pm\infty$

$$f(\xi) \rightarrow 2\pi n_{\pm}, \quad f'(\xi) \rightarrow 0 \quad (\Rightarrow f''(\xi) \rightarrow 0),$$

where the two integers  $n_{\pm} \in \mathbb{Z}$  can be different. Indeed they are different for a kink (/antikink) solution. Choosing the branch of the arctan such that

$$\arctan(0^{\pm}) = 0^{\pm}, \quad \arctan(\pm\infty) = \pm \left(\frac{\pi}{2}\right)^{\mp},$$

we find that the kink and the anti-kink solution look as in fig. 2.2 at a fixed time  $t$ :




Figure 2.2: Snapshots of the sine-Gordon kink and anti-kink.

### REMARKS:

1. Choosing a different branch of the  $\arctan$ <sup>3</sup> shifts the whole solution  $u(x, t)$  by a multiple of  $2\pi$ . This is inconsequential. What matters is:

$$\begin{aligned} u(+\infty, t) - u(-\infty, t) &= +2\pi && \text{KINK} \\ u(+\infty, t) - u(-\infty, t) &= -2\pi && \text{ANTI-KINK} \end{aligned}$$

2. The velocity of the kink/anti-kink could be

$$\begin{aligned} v > 0 : & \quad \text{RIGHT-MOVING} \\ v = 0 : & \quad \text{STATIC} \\ v < 0 : & \quad \text{LEFT-MOVING} \end{aligned}$$

3. For a **real** solution we need

$$\gamma^2 \geq 0 \quad \implies \quad |v| \leq 1 = \text{speed of light}$$

4. The kink/antikink is a **localised** lump centred at  $x_0 + vt$  and with

$$\text{WIDTH} \sim \frac{1}{\gamma} = \sqrt{1 - v^2}.$$

<sup>3</sup>along with reversing the sign and adjusting the integration constant if the multiple is odd. Check for yourself.

So faster kinks/antikinks are narrower. This phenomenon is known as “Lorentz contraction” and is a feature of special relativity.  $\gamma$  is called the “**Lorentz factor**”.

**NOTE:** It might be confusing to state that the kink/antikink is localised, when  $u$  interpolates between different values as  $x \rightarrow \pm\infty$ . The key point is that  $u$  is an angular variable which is only defined modulo addition of  $2\pi$ . To define the width it is better to look at single-valued objects like  $e^{iu}$  or  $\partial_x u$ , which do not suffer from the above ambiguity. This point will become more concrete later when we calculate the energy density of the kink, which is a single-valued and everywhere positive function, which achieves a maximum at the centre of the kink and approaches zero far away from the centre, see figure 3.2.

## 2.3 A mechanical model for the sine-Gordon equation

Consider a chain of infinitely many identical pendula hanging from a straight wire which cannot be stretched but can be twisted. Each identical pendulum consists of a massless<sup>4</sup> rod of length  $L$ , with a weight of mass  $M$  at the end of the rod. The pivot of the  $n$ -th pendulum is at position  $na$  along the line, where  $n \in \mathbb{Z}$  and  $a$  is the separation, and the configuration of the  $n$ -th pendulum at time  $t$  is encoded by  $\theta_n(t)$ , the angle between the pendulum and the downward pointing vertical at time  $t$ . See figure 2.3.




Figure 2.3: Section of an infinite chain of pendula separated by distance  $a$ .

The pendula are subject to two kinds of forces: a gravitational force due to the attraction between the Earth and the weights, which favours downward pointing pendula; and a twisting force between neighbouring pendula due to the wire, which favours a straight untwisted wire and therefore the alignment of neighbouring pendula.<sup>5</sup> The **equations of motion** (the

<sup>4</sup>This assumption can be easily relaxed, leading to no qualitative difference in what follows.

<sup>5</sup>This is a slight lie. If you have studied rigid bodies you will recognise that these are “torques” rather than forces. The equation of motion (2.3) is not the standard Newton’s law  $F = ma$ , but rather its rotational analogue, which states that the total torque equals the product of the moment of inertia and the angular acceleration.

analogue of Newton's equation  $F = ma$ ) for this physical system are a coupled system of infinitely many ODE's labelled by the integer  $n$ , one for each pendulum, which take the form

$$ML^2\ddot{\theta}_n(t) = \underbrace{-MgL \cdot \sin \theta_n(t)}_{\text{net gravitational force}} + \underbrace{\frac{k}{a}(\theta_{n+1}(t) - \theta_n(t)) + \frac{k}{a}(\theta_{n-1}(t) - \theta_n(t))}_{\text{twisting forces exerted by neighbouring pendula}}, \quad n \in \mathbb{Z} \quad (2.3)$$

where a dot denotes a time derivative,  $g$  is the gravitational acceleration and  $k$  is an elastic constant that parametrizes the strength of the twisting force.

Now we are going to take the so called “**continuum limit**” of this infinite-dimensional discrete system, in which the separation between consecutive pendula becomes infinitesimally small and the average mass density (*i.e.* the mass per unit length) along the line is kept fixed:

$$a \rightarrow 0, \quad m = M/a \text{ fixed}.$$

In the continuum limit, the position  $x = na$  of the  $n$ -th pendulum along the line effectively becomes a continuous real variable, which replaces the discrete index  $n \in \mathbb{Z}$ . Identifying  $\theta_n(t) \equiv \theta(x = na, t)$ , the collection  $\{\theta_n(t)\}_{n \in \mathbb{Z}}$  of angular coordinates of the infinitely many pendula at time  $t$  is replaced in the limit by a single function  $\theta(x, t)$  of two continuous variables, space and time. By the definition of the derivative as a limit, we also have that

$$\begin{aligned} \frac{\theta_{n+1}(t) - \theta_n(t)}{a} &\rightarrow \theta'(x, t), \\ \frac{1}{a} \left( \frac{\theta_{n+1}(t) - \theta_n(t)}{a} - \frac{\theta_n(t) - \theta_{n-1}(t)}{a} \right) &\rightarrow \theta''(x, t). \end{aligned}$$

where a prime denotes an  $x$ -derivative.

Dividing the equations of motion (2.3) by  $ML^2 = amL^2$  and taking the continuum limit we find the single equation of motion

$$\ddot{\theta} = -\frac{g}{L} \sin \theta + \frac{k}{mL^2} \theta''$$

for the “**field**”  $\theta(x, t)$ . We can get rid of the constants by rescaling  $x$  and  $t$ <sup>6</sup>, and rearrange to get the equation

$$\ddot{\theta} - \theta'' = -\sin \theta,$$

which is nothing but the sine-Gordon equation  $\theta_{tt} - \theta_{xx} = -\sin \theta$  for the field  $\theta$ ! We say therefore that the sine-Gordon equation is the continuum limit of (2.3).

We can use this mechanical model to gain some intuition about the possible configurations of the sine-Gordon field:

---

<sup>6</sup>Send  $x \mapsto \sqrt{\frac{k}{mgL}} x$  and  $t \mapsto \sqrt{\frac{L}{g}} t$ .

- The **lowest energy state** (or “**ground state**”, or “**vacuum**”) of the system is the configuration with all pendula pointing downwards,

$$\theta(x, t) = 0 \pmod{2\pi} \quad \forall x ,$$

which is a configuration of stable equilibrium.<sup>7</sup> See figure 2.4.




Figure 2.4: Configuration of stable equilibrium for the chain of pendula.

- By a continuous perturbation of the vacuum, we can obtain configuration which represents a “small wave”, which satisfies the same boundary conditions of the vacuum,  $\theta \rightarrow 0$  as  $x \rightarrow \pm\infty$ .<sup>8</sup>




Figure 2.5: A small wave going through the chain of pendula.

- There are also configurations in which the chain of pendula twists around the line. If they twist once in the direction of increasing angles, so that  $\theta$  increases by  $2\pi$  from  $x \rightarrow -\infty$  to  $x \rightarrow +\infty$ , this describes a kink or a continuous deformation thereof:

If instead they twist once in the direction of decreasing angles, so that  $\theta$  decreases by  $2\pi$  from  $x \rightarrow -\infty$  to  $x \rightarrow +\infty$ , this describes an anti-kink or a continuous deformation thereof.

- The **limiting values** of the sine-Gordon field  $\theta$  as  $x \rightarrow \pm\infty$  are **fixed**: changing them would require twisting infinitely many pendula by 360 degrees, which would cost energy.

<sup>7</sup>We will confirm this intuition later when we study the energy of the sine-Gordon field.

<sup>8</sup>We will see later that this “small wave” does not need to be small, in fact. For instance it could look like a kink followed by an antikink.






Figure 2.6: A kink going through the chain of pendula.

If

$$\theta(+\infty, t) - \theta(-\infty, t) = 2m\pi, \quad \text{with } m \neq 0 \text{ integer},$$

then the configuration of the system **cannot be deformed continuously to the vacuum** where all pendula point downwards, unlike the “small wave” mentioned above. This tells us that the kink (or the antikink) **cannot disperse/dissipate into the vacuum**. This is related to the notion of **topological stability**, which we will discuss in the next chapter.

I invite you to play with this Wolfram demonstration of the chain of coupled pendula, using Mathematica (which should be available on university computers – let me know if it isn’t) or the free Wolfram Player. Play with the parameters and visualise a kink, the scattering of two kinks or of a kink and an anti-kink, and the breather, a bound state of a kink and an anti-kink. We will study all of these configurations in the continuum limit later in the term, using the sine-Gordon equation.

## 2.4 Travelling waves and 1d point particles

Looking for a travelling wave solutions  $u(x, t) = f(x - vt) \equiv f(\xi)$  of the KdV and sine-Gordon equation, we encountered equations of the form

$$f'' = \hat{F}(f)$$

where a prime denotes a derivative with respect to  $\xi$ . We integrated this equation to

$$\frac{1}{2}(f')^2 + \hat{V}(f) = \hat{E} = \text{const} \quad (*)$$

where

$$\hat{V}(f) = - \int df \hat{F}(f).$$




Figure 2.7: Example of a potential energy  $V(x)$  and force  $F(x) = -V'(x)$ .

By tuning the integration constant in this indefinite integral and absorbing it in  $\hat{E}$ , we can set  $\hat{E}$  to zero or to any value we wish.

The previous equations are **analogous** to the **classical mechanics** of a **point particle** moving in **one space dimension**. Let  $x(t)$  be the position of the point particle at time  $t$  and dots denote time derivatives. The equation of motion (EoM) of the point particle is **Newton's equation**

$$m\ddot{x} = F(x)$$

(mass  $\times$  acceleration = force) can be integrated to the **energy conservation** law

$$\frac{1}{2}m\dot{x}^2 + V(x) = E = \text{const}$$

(kinetic energy + potential energy = total energy, which is constant in time), where the force and the potential energy are related by

$$F(x) = -\frac{d}{dx}V(x).$$

The potential energy and the total energy can be shifted by a common constant with no physical change. See figure 2.7 for an example of a potential energy  $V(x)$  and the associated force  $F(x) = -V'(x)$ .

It may be useful to think of  $x$  as the horizontal coordinate of a point particle (think of an infinitesimal ball) moving on a hill of vertical height  $V(x)$  at coordinate  $x$ , subject only to the gravitational force and the reaction of the ground (which is equal and opposite when the ground is flat). Even if you are not very familiar with classical mechanics, you will hopefully have some intuition of what will happen to the ball.<sup>9</sup>

<sup>9</sup>You can also model this by riding a brakeless bike in hilly Durham. It's a good idea to develop some intuition about this physical system without running the experiment yourself, which I don't recommend. (This is one of a number of reasons why theoretical physics is superior to experimental physics.)

The mathematical correspondence between the equations for a travelling wave in one space and one time dimension and for a classical point particle in one space dimension is

$\xi$	$\longleftrightarrow$	$t$
$f$	$\longleftrightarrow$	$x$
1	$\longleftrightarrow$	$m$
$\hat{F}(f)$	$\longleftrightarrow$	$F(x)$
$\hat{E} - \hat{V}(f)$	$\longleftrightarrow$	$E - V(x)$

This correspondence allows us to understand the qualitative behaviour of travelling waves even when we cannot integrate equation (\*) exactly, using elementary facts from classical mechanics, which are encoded in the the mathematics of the previous equations:


1. The total **energy is conserved** and can only be converted from kinetic energy (which is non-negative!) to potential energy and vice versa. The velocity  $\dot{x}$  of the point particle is zero if and only if the kinetic energy is zero, which means that all the energy is stored in potential energy:

$$\dot{x} = 0 \quad \Longleftrightarrow \quad V(x) = E .$$

2. When the point particle reaches one of the special values of  $x$  such that  $V(x) = E$ , either of two things happens depending on the acceleration of the particle:

(a)  $F(x) = -\frac{d}{dx}V(x) \neq 0$ :


The acceleration is non-vanishing, therefore the particle **reverses its direction of motion**:



These values of  $x$  are known as “**turning points**”.

(b)  $F(x) = -\frac{d}{dx}V(x) = 0$ :

The acceleration vanishes and the particle **stops**.




These values of  $x$  are known as “**equilibrium points**”. The approach to equilibrium takes an infinite time.

**\* EXERCISE:** Derive the previous statements by Taylor expanding the potential energy about a point where  $V(x) = E$  and substituting the expansion in the energy conservation law.

Now let us translate this discussion to the context of travelling waves. We will focus on the examples of the KdV and the sine-Gordon equation here, but more examples are available in **[Ex 13]** in the problems set.

### EXAMPLES:


1. **KdV:**  $\hat{E} = 0$ ,  $\hat{V}(f) = f^2 \left( f - \frac{v}{2} \right)$  ( $v > 0$ )



From a graphical analysis of  $\hat{V}(f)$  and the analogy between travelling waves and point particles in one dimension, we see that there exists a travelling wave solution that starts at  $f = 0^+$  at  $\xi \rightarrow -\infty$ , increases until the ‘turning point’  $f = v/2$ , and decreases to  $f = 0^+$  at  $\xi \rightarrow +\infty$ . This is nothing but the KdV soliton (2.1) that we found in section 2.1. If instead the travelling wave solution starts at  $f = 0^-$  at  $\xi \rightarrow -\infty$ , then it will fall down the cliff and reach  $f \rightarrow -\infty$ , leading to a singular solution, that we discard. Note that if  $v < 0$  we have that  $\hat{V}(0) = 0$ , but  $\hat{V}(f) > 0$  for small  $f \neq 0$ . Therefore the only

real solution obeying the boundary conditions is the constant zero solution  $f(\xi) = 0$  for all  $\xi$ . If  $v = 0$ , in addition to the trivial solution there is also a singular real travelling wave solution that we discard on physical grounds.

2. **sine-Gordon:**  $\hat{E} = 0$ ,  $\hat{V}(f) = \gamma^2 (\cos f - 1)$



From a graphical analysis of  $\hat{V}(f)$ , we see that two classes of travelling wave solutions exist: one where  $f$  interpolates between  $2n\pi$  at  $x \rightarrow -\infty$  and  $2(n+1)\pi$  at  $x \rightarrow \infty$ , and another where  $f$  interpolates between  $2n\pi$  at  $x \rightarrow -\infty$  and  $2(n-1)\pi$  at  $x \rightarrow \infty$ . We identify these solutions with the kink and anti-kink (2.2) of section 2.2.

**\* EXERCISE:** Using the analogy with a one-dimensional point particle, determine the qualitative behaviour of a travelling wave solution of the KdV equation on a circle (i.e. with periodic boundary conditions). [**Hint:** allow integration constants  $A, B \neq 0$  and look at  $\hat{V}(f)$ .] **[Ex 14\*]**

# Chapter 3

## Topological lumps and the Bogomol'nyi bound

The main references for this chapter are §5.3, 5.1 of [Manton and Sutcliffe, 2004] and §2.1 of [Dauxois and Peyrard, 2006].

### 3.1 The sine-Gordon kink as a topological lump

In chapter 2 I mentioned the **topological properties** of the **sine-Gordon kink**, which ensure that it cannot disperse or dissipate to the vacuum. Let us understand these topological properties better. As a reminder, the sine-Gordon equation for the field  $u$  is

$$u_{tt} - u_{xx} + \sin u = 0 .$$

Starting from the **discrete** mechanical model involving pendula of section 2.3, rescaling  $x$  and  $t$  as in footnote 6 so as to eliminate all constants, and taking the continuum limit  $a \rightarrow 0$ , it is not hard to see that the **kinetic energy**  $T$  and the **potential energy**  $V$  of the sine-Gordon field are **[Ex 15]**

$$T = \int_{-\infty}^{+\infty} dx \frac{1}{2} u_t^2 \tag{3.1}$$

$$V = \int_{-\infty}^{+\infty} dx \left[ \underbrace{\frac{1}{2} u_x^2}_{\text{twisting}} + \underbrace{(1 - \cos u)}_{\text{gravity}} \right] . \tag{3.2}$$

#### **REMARK:**

The kinetic and potential energies of the sine-Gordon field are the continuum limits of the

kinetic and potential energies of the infinite chain of pendula. They should not be confused with  $\frac{1}{2}(f')^2$  and  $\hat{V}(f)$  for the one-dimensional point particle in the analogy of section 2.4.

We can use this result to deduce the boundary conditions that we anticipated in section 2.2. The boundary conditions follow from requiring that **all field configurations** have **finite (total) energy**  $E = T + V$ . Since the total energy is the integral over the real line of the sum of three non-negative terms, all three terms must vanish in the asymptotic limits  $x \rightarrow \pm\infty$  to ensure the convergence of the integral. So the finiteness of the energy requires the boundary conditions

$$u_t, \quad u_x, \quad 1 - \cos u \quad \xrightarrow{x \rightarrow \pm\infty} \quad 0 \quad \forall t.$$

Since  $1 - \cos u = 0$  iff  $u$  is an integer multiple of  $2\pi$ , we need

$$\boxed{u(-\infty, t) = 2\pi n_-, \quad u(+\infty, t) = 2\pi n_+,} \quad (3.3)$$

for some integers  $n_{\pm}$ . (This means that pendula are at rest, pointing downwards, as  $x \rightarrow \pm\infty$ .)

### **REMARKS:**<sup>1</sup>

1. The overall value of  $n_{\pm}$  has no meaning, since  $u$  is defined modulo  $2\pi$ . A shift of the field  $u \mapsto u + 2\pi k$  is unphysical, but it shifts  $n_{\pm} \mapsto n_{\pm} + k$ . What really matters is the difference  $n_+ - n_-$ , which is invariant under this ambiguity:

$$\frac{1}{2\pi} [u(+\infty, t) - u(-\infty, t)] = n_+ - n_- = \# \text{ of "twists"/"kinks"}$$

2. The integer  $n_+ - n_-$  is "**TOPOLOGICAL**", *i.e.* it does not change under any **continuous** changes of the field  $u$  (and of the energy  $E$ ). In particular, it cannot change under time evolution, since time is continuous. Therefore it is a **constant of motion** or a "**conserved charge**" (more about this in the next chapter). Since the conservation of  $n_+ - n_-$  is due to a topological property, we call this a "**TOPOLOGICAL CHARGE**".<sup>2</sup> Solutions with the same topological charge are said to belong to the same "**TOPOLOGICAL SECTOR**".

---

<sup>1</sup>Some of these remarks were made for kinks and antikinks in the previous chapter. Now that we derive them from the BC's, we see that they hold more generally for all solutions.

<sup>2</sup>[Advanced remark for those who know some topology – if you don't, you can safely ignore this:] Mathematically,  $n_+ - n_-$  is a "winding number", the topological invariant which characterises maps  $S^1 \rightarrow S^1$ . The first  $S^1$  is the compactification of the spatial real line, with the points at infinity identified, and the second  $S^1$  is the circle parametrised by  $u \bmod 2\pi$ . The winding number counts how many times  $u$  winds around the circle as  $x$  goes from  $-\infty$  to  $+\infty$ .

3. Dispersion and dissipation occur by time evolution, a continuous process which cannot change the value of the integer  $n_+ - n_-$ . Since the vacuum has  $n_+ - n_- = 0$ , any configuration with  $n_+ - n_- \neq 0$  **cannot disperse/dissipate to the vacuum**.

### VOCABULARY:

- “**TOPOLOGICAL CONSERVATION LAW**”:

The conservation (in time) of a topological charge, that is  $\frac{d}{dt}(\text{topological charge}) = 0$ .

- “**TOPOLOGICAL LUMP**”:

A localised field configuration which cannot dissipate or disperse to the vacuum by virtue of a topological conservation law.

So the **sine-Gordon kink** is a **topological lump**. It is also a **soliton**, but to see that we will have to check property 3, which concerns scattering.

Topological lumps also exist in higher dimensions. A notable example is the “magnetic monopole”, a magnetically charged localised object that exists in certain generalizations of electromagnetism in three space and one time dimensions. Another example is the “vortex”, which is a topological lump if space is  $\mathbb{R}^2$ .<sup>3</sup>

## 3.2 The Bogomol'nyi bound

Among the kink solutions found in (2.2) using the travelling wave *ansatz*, there was a **STATIC KINK** with zero velocity. Topology tells us that it cannot disperse or dissipate completely to the vacuum. But **is its precise shape “stable”** under small perturbations? This would be **guaranteed if** we could show that **it minimises the energy** amongst all configurations with the same topological charge. The reason is that any perturbation near a minimum of the energy would increase the energy, which however is conserved upon time evolution.<sup>4</sup>

A useful analogy to keep in mind is with a point particle on a hilly landscape under the force of gravity, as in figure 3.1: if the point particle is sitting still at a local minimum of the height, minimising the energy (locally), it is in a configuration of stable equilibrium. Any perturbation would necessarily move the particle up the hill, but this is not allowed under time evolution as it would increase the total energy.

---

<sup>3</sup>Indeed there is a topological charge, the ‘vortex number’, which is conserved and can be non-vanishing if space is  $\mathbb{R}^2$ . On the other hand, topology implies that the vortex number vanishes on the two-sphere  $S^2$ : this is fortunate, because if it were non-vanishing there would always be hurricanes going around the surface of Earth.

<sup>4</sup>We will in fact show that the static kink is a global minimum of the energy amongst configurations with unit topological charge. This ensures its stability even when one includes quantum effects, which we are not






Figure 3.1: A point particle on a hilly landscape is stable if it locally minimises the energy. This happens when it is sitting still at a minimum of the potential energy.

So we will seek a lower bound for the total energy  $E = T + V$  in the topological sector of the kink, which has topological charge  $n_+ - n_- = 1$ . The energy is the integral of a non-negative energy density, so immediately find the lower bound  $E \geq 0$ , but we can do better than that:

$$\begin{aligned}
 E = T + V &= \int_{-\infty}^{+\infty} dx \left[ \frac{1}{2} u_t^2 + \frac{1}{2} u_x^2 + (1 - \cos u) \right] \\
 &\stackrel{(u_t^2 \geq 0)}{\geq} \int_{-\infty}^{+\infty} dx \left[ \frac{1}{2} u_x^2 + (1 - \cos u) \right] \\
 &= \int_{-\infty}^{+\infty} dx \left[ \frac{1}{2} u_x^2 + 2 \sin^2 \frac{u}{2} \right] \\
 &\stackrel{\text{"Bogomol'nyi trick"}}{=} \int_{-\infty}^{+\infty} dx \left[ \frac{1}{2} \left( u_x \pm 2 \sin \frac{u}{2} \right)^2 \mp 2 \sin \frac{u}{2} \cdot u_x \right] \\
 &= \int_{-\infty}^{+\infty} dx \frac{1}{2} \left( u_x \pm 2 \sin \frac{u}{2} \right)^2 \pm 4 \left[ \cos \frac{u}{2} \right]_{-\infty}^{+\infty}. \quad (*)
 \end{aligned}$$

A few comments are in order:

1. The inequality in the second line follows from omitting the non-negative term  $\frac{1}{2} u_t^2$ . It is **"saturated"** (that is, it becomes an equality) for **static field configurations**, such that  $u_t = 0$ ;
2. In the third line we used a half-angle formula;
3. In the fourth line we used the so called **"Bogomol'nyi trick"** to replace a sum of squares by the square of a sum plus a correction term which is a total  $x$ -derivative;
4. In the fifth line we integrated the total derivative, leading to a **"boundary term"** (or **"surface term"**) which only depends on the limiting values of the field at spatial infinity.

---

concerned with in this course.

If  $u$  satisfies the 1-**kink BC's**

$$\boxed{u(-\infty, t) = 0, \quad u(+\infty, t) = 2\pi},$$

then the boundary term evaluates to

$$4 \left[ \cos \frac{u}{2} \right]_{-\infty}^{+\infty} = 4(-1 - 1) = -8.$$

Picking the lower (*i.e.*  $-$ ) signs in (\*), we obtain the lower bound

$$\boxed{E \geq \int_{-\infty}^{+\infty} dx \frac{1}{2} \left( u_x - 2 \sin \frac{u}{2} \right)^2 + 8 \geq 8} \quad (3.4)$$

for the energy, where the second inequality is saturated if the expression in brackets vanishes.<sup>5</sup> Equation (3.4) is known as the “**BOGOMOL'NYI BOUND**”.

The **Bogomol'nyi bound** (3.4) is **saturated** (*i.e.*  $E = 8$ ) if and only if the field configuration is **static**, that is

$$\boxed{u_t = 0},$$

and satisfies the “**BOGOMOL'NYI EQUATION**”

$$\boxed{u_x = 2 \sin \frac{u}{2}}. \quad (3.5)$$

So we can find the **least energy field configurations** in the “**1-kink topological sector**” (*i.e.* with  $n_+ - n_- = 1$ ) by looking for static solutions  $u = u(x)$  of the Bogomol'nyi equation:

$$u_x = 2 \sin \frac{u}{2} \implies \int dx = \int \frac{du}{2 \sin \frac{u}{2}} = \log \tan \frac{u}{4},$$

whose general solution is

$$\boxed{u(x) = 4 \arctan \left( e^{x-x_0} \right)}. \quad (3.6)$$

This is nothing but the **static kink**, which we obtained in section 2.2 as a special case of a travelling wave solution of the sine-Gordon equation with  $v = 0$ .

#### **REMARK:**

Note that the Bogomol'nyi equation, being a first order differential equation (in fact an ODE once we impose  $u_t = 0$ ), is much easier to solve than the full equation of motion, the sine-Gordon equation, which is a second order PDE.

---

<sup>5</sup>Picking the upper (*i.e.*  $+$ ) signs in (\*) we obtain the lower bound  $E \geq -8$ , which is weaker than the trivial bound  $E \geq 0$  therefore not very useful. The Bogomol'nyi trick always has a sign ambiguity. The choice of sign that leads to the stricter inequality depends on the sign of the boundary term.




Figure 3.2: The energy density of a static kink.

**\* EXERCISE:** Check that a field configuration that saturates the Bogomol'nyi bound is automatically a solution of the sine-Gordon equation.

So we learned that **amongst all solutions with topological charge  $n_+ - n_- = 1$ , the static kink has the least energy**, hence it is **stable**. Indeed, topology in principle allows the kink to disperse to other solutions with  $n_+ - n_- = 1$ , but the dispersing waves would carry some of the energy away. Since the static kink has the least energy in the  $n_+ - n_- = 1$  topological sector, it can't lose energy, hence it's stable. This notion of stability which originates from minimising the energy in a given topological sector is called **"TOPOLOGICAL STABILITY"**.

Using staticity and the Bogomol'nyi equation, we now have a shortcut to compute the **energy density  $\mathcal{E}$**  of the **static kink**, namely the integrand of the total energy  $E = \int_{-\infty}^{+\infty} dx \mathcal{E}$ :

$$\mathcal{E} = \frac{1}{2}u_t^2 + \frac{1}{2}u_x^2 + 2\sin^2 \frac{u}{2} \underset{\substack{u_t=0 \\ u_x=2\sin \frac{u}{2}}}{=} u_x^2 = 4\text{sech}^2(x - x_0),$$

which shows that the energy density of the kink is localised near  $x_0$ , see figure 3.2.

**\* EXERCISE:** Think about how to generalise the Bogomol'nyi bound for higher topological charge, for instance  $n_+ - n_- = 2$ . This is not obvious! **[Ex 17]**


### 3.3 Summary

There are two ways for a lump to be long-lived:

1. by **INTEGRABILITY** (infinitely many conservation laws, more about this next)  
→ **"TRUE" (or "INTEGRABLE") SOLITONS**
2. by **TOPOLOGY** (topological conservation law)

→ **TOPOLOGICAL LUMPS.**<sup>6</sup>

It is important to note that these two mechanisms are not mutually exclusive: there are some lumps, like the sine-Gordon kink, which are both topological lumps and true solitons. The various possibilities and some examples are summarised in the following Venn diagram:



---

<sup>6</sup>Some people use the term solitons for both integrable solitons and topological lumps, but in this course we will only refer to the former as “solitons”).

# Chapter 4

## Conservation laws

The main references for this chapter are §5.1.1 and §5.1.2 of [Drazin and Johnson, 1989].

**Conservation laws** provide the most fundamental characterisation of a physical system: they tell us which quantities don't change with time. For the purpose of this course, they play a key role because they explain why the motion of “true” solitons is so restricted that they scatter without changing their shapes.

The idea of a conservation law is to construct spatial integrals of functions of the field  $u$  and its derivatives

$$Q = \int_{-\infty}^{+\infty} dx \, \rho(u, u_x, u_{xx}, \dots, u_t, u_{tt}, \dots) \quad (4.1)$$

which are constant in time (in physics parlance, they are “**constants of motion**”)

$$\frac{d}{dt} Q = 0 \quad (4.2)$$

when  $u$  satisfies its **equation of motion (EoM)**, such as the sine-Gordon equation or the KdV equation. The constant of motion (4.1) is called a “**CONSERVED CHARGE**” or “**CONSERVED QUANTITY**” and the equation (4.2) stating its time-independence is called a “**CONSERVATION LAW**”.

For the KdV and the sine-Gordon equation, it turns out that there exist **infinitely many conserved quantities**. This makes them “**integrable systems**” (more about this next term) and explains many of their special properties.

## 4.1 The basic idea

The standard method for constructing a conserved charge like (4.1) involves finding two functions  $\rho$  and  $j$  of  $u$  and its derivatives, such that the EoM for  $u$  implies the “**LOCAL CONSERVATION LAW**” or “**CONTINUITY EQUATION**”

$$\boxed{\frac{\partial \rho}{\partial t} + \frac{\partial j}{\partial x} = 0} \quad (4.3)$$

and the boundary conditions imply<sup>1</sup>

$$\boxed{\lim_{x \rightarrow +\infty} j - \lim_{x \rightarrow -\infty} j = 0 \quad \forall t.} \quad (4.4)$$

Then

$$\frac{d}{dt} \int_{-\infty}^{+\infty} dx \rho = \int_{-\infty}^{+\infty} dx \frac{\partial \rho}{\partial t} \stackrel{(4.3)}{=} - \int_{-\infty}^{+\infty} dx \frac{\partial j}{\partial x} = -[j]_{-\infty}^{+\infty} \stackrel{(4.4)}{=} 0.$$

Hence

$$Q = \int_{-\infty}^{+\infty} dx \rho \quad (4.5)$$

is a **conserved CHARGE**.  $\rho$  is called the conserved “**CHARGE DENSITY**”, and  $j$  is called the conserved “**CURRENT DENSITY**” (or just “**CURRENT**”, by a common abuse of terminology.)

## 4.2 Example: conservation of energy for sine-Gordon

Is the total energy

$$\boxed{E = \int_{-\infty}^{+\infty} dx \mathcal{E}}$$


conserved for the sine-Gordon field, where the energy density is

$$\boxed{\mathcal{E} = \frac{1}{2}u_t^2 + \frac{1}{2}u_x^2 + (1 - \cos u)} \quad ? \quad (4.6)$$

The energy density  $\mathcal{E}$  plays the role of  $\rho$  here. Can we show then that  $\rho = \mathcal{E}$  obeys a continuity equation (4.3) for some function  $j$  that obeys the limit condition (4.4), when the sine-Gordon equation (EoM)

$$u_{tt} - u_{xx} + \sin u = 0$$

---

<sup>1</sup>  Don't forget that  $\infty - \infty$  is not equal to zero: it's undetermined. So in particular  $j$  must have finite limits at spatial infinity for (4.4) to hold.

holds? Let's compute:

$$\begin{aligned}
 \frac{\partial \mathcal{E}}{\partial t} &= u_t u_{tt} + u_x u_{xt} + \sin u \cdot u_t \\
 &= u_t (u_{tt} + \sin u) + u_x u_{xt} \\
 &\stackrel{\text{EoM}}{=} u_t u_{xx} + u_x u_{xt} = \frac{\partial}{\partial x} (u_t u_x) \equiv \frac{\partial}{\partial x} (-j) ,
 \end{aligned}$$

and since the BC's for the sine-Gordon equation imply that  $u_t u_x \rightarrow 0$  as  $x \rightarrow \pm\infty$ , we deduce that **energy is conserved**:

$$\boxed{\frac{dE}{dt} = 0} .$$

### 4.3 Extra conservation laws for relativistic field equations

For any **relativistic field theory** in 1 space ( $x$ ) + 1 time ( $t$ ) dimensions (e.g. Klein-Gordon, sine-Gordon, “ $\phi^4$ ”, ...), the **energy**

$$\boxed{E = \int_{-\infty}^{+\infty} dx \mathcal{E} = \int_{-\infty}^{+\infty} dx \left[ \frac{1}{2} u_t^2 + \frac{1}{2} u_x^2 + V(u) \right]} \quad (4.7)$$

is **conserved**, when the equation of motion (**EoM**)

$$\boxed{u_{tt} - u_{xx} = -V'(u)} \quad (4.8)$$

is **satisfied**.

**\* EXERCISE:** Check this statement.

The “**scalar potential**”  $V(u)$  determines the theory. For instance

$$V(u) = \begin{cases} \frac{1}{2} m^2 u^2 & \text{(Klein-Gordon)} \\ 1 - \cos u & \text{(sine-Gordon)} \\ \frac{\lambda}{2} (u^2 - a^2)^2 & \text{ (“}\phi^4\text{”)} \\ \dots & \end{cases}$$

A deep theorem due to Emmy Noether shows that the **conservation of energy** follows from the **invariance** of the theory **under arbitrary time translations**  $t \mapsto t + c$ . Similarly, **invariance under space translations**  $x \mapsto x + c'$  implies the **conservation of momentum**  $P$ .

We will not delve into Noether's theorem, as you might encounter it in other courses and also because it is of limited help for our purposes. The **question** that we would like to answer now is:

**Can there be more conservation laws, in addition to energy and momentum conservation?**

We will answer this question constructively.

The **first step** is to switch to **light-cone coordinates**

$$\boxed{x^\pm = \frac{1}{2}(t \pm x)} \iff \begin{cases} t &= x^+ + x^- \\ x &= x^+ - x^- \end{cases}, \quad (4.9)$$

which are so called because the trajectory of light rays is  $x^+ = \text{const}$  or  $x^- = \text{const}$  for left-moving or right-moving rays respectively. By the chain rule we calculate

$$\begin{aligned} \partial_\pm &\equiv \frac{\partial}{\partial x^\pm} = \frac{\partial t}{\partial x^\pm} \frac{\partial}{\partial t} + \frac{\partial x}{\partial x^\pm} \frac{\partial}{\partial x} = \frac{\partial}{\partial t} \pm \frac{\partial}{\partial x} \equiv \partial_t \pm \partial_x \\ &\implies \partial_+ \partial_- = \partial_t^2 - \partial_x^2, \end{aligned}$$

so the EoM can be written synthetically as

$$\boxed{u_{+-} = -V'(u)}, \quad (4.10)$$

where we used the shorthand notation  $f_\pm \equiv \frac{\partial f}{\partial x^\pm} \equiv \partial_\pm f$ .

Next, we define the “**Lorentz spin**” of a monomial built out of light-cone derivatives of functions of  $u$  as the **number of  $\partial_+$  derivatives minus the number of  $\partial_-$  derivatives**. For instance  $(u_+)^3 u_- u_{+-}$  has spin  $3 - 1 + (2 - 1) = 3$ . According to the theory of special relativity, objects of different spins transform differently under the “Lorentz group” of symmetries of relativistic field equations. If you would like to know more about Lorentz transformations and the Lorentz spin, you can read this optional note. If you don't know any relativity and don't want to read that note, you'll just need to remember that objects of different spins don't talk to one another, and that it only makes sense to add/subtract objects of the same spin. Note in particular that we can use the EoM (4.10) to convert  $u_{+-}$  into  $-V'(u)$ , which does not affect the spin. So from now on we will focus on objects of definite spin, and we will treat different spins separately.

Next, suppose that there exists a **pair of densities**  $(T_{s+1}, X_{s-1})$  of spins  $s \pm 1$  such that

$$\boxed{\partial_- T_{s+1} = \partial_+ X_{s-1}}. \quad (4.11)$$



In terms of the original space and time coordinates  $x$  and  $t$ , this is nothing but the continuity equation<sup>2</sup>

$$\partial_t \underbrace{(T_{s+1} - X_{s-1})}_{\rho} + \partial_x \underbrace{(-T_{s+1} - X_{s-1})}_{j} = 0 .$$

Therefore, by the basic idea of section 4.1, provided that

$$\lim_{x \rightarrow +\infty} (T_{s+1} + X_{s-1}) - \lim_{x \rightarrow -\infty} (T_{s+1} + X_{s-1}) = 0 , \quad (4.12)$$

it follows that

$$Q_s = \int_{-\infty}^{+\infty} dx (T_{s+1} - X_{s-1}) \quad (4.13)$$

is a **conserved charge** (which turns out to have spin  $s$ , hence the subscript).

In order to construct conserved charges (4.13), our aim will be to find  $(T, X)$  pairs which satisfy equations (4.11)-(4.12). We will make a **simplifying assumption**: we will only consider “**polynomially conserved densities**”

$$T = (\text{polynomial in } x^+ \text{-derivatives of } u) ,$$

so we will never consider  $u$  without derivatives, and  $T_{s+1}$  will be a sum of terms with precisely  $s + 1$  derivatives  $\partial_+$  and no  $\partial_-$  derivatives.

We will also disregard total  $x^+$ -derivatives in  $T$  and consider  $(T, X)$  pairs modulo the equivalence relation

$$(T_{s+1}, X_{s-1}) \underset{\text{“equivalent to”}}{\sim} (T'_{s+1}, X'_{s-1}) = (T_{s+1} + \partial_+ U_s, X_{s-1} + \partial_- U_s) , \quad (4.14)$$

so long as

$$\lim_{x \rightarrow +\infty} U_s - \lim_{x \rightarrow -\infty} U_s = 0 \quad \forall t . \quad (4.15)$$

This is because the shifts in (4.14) cancel out in the continuity equation (4.11) and in the charge:

$$Q'_s = \int_{-\infty}^{+\infty} dx (T'_{s+1} - X'_{s-1}) \underset{(4.14)}{=} Q_s + \int_{-\infty}^{+\infty} dx (\partial_+ - \partial_-) U_s = Q_s + 2 \int_{-\infty}^{+\infty} dx \partial_x U_s \underset{(4.15)}{=} Q_s .$$

Let us proceed spin by spin.

---

<sup>2</sup>The reason why densities of different spins are mixed in the next equation, seemingly contradicting the statement I have made a few lines above, is that  $\partial_t$  and  $\partial_x$  don't have definite spin either. In (4.11), which uses light-cone derivatives  $\partial_{\pm}$ , both sides have the same spin  $s$ , consistently with my statement.

$$\boxed{s = 0} \quad T_1 = u_+$$

is the only polynomially conserved density of spin 1, up to an irrelevant multiplicative factor which can be absorbed in the normalisation of the charge. It solves (4.11) with  $X_{-1} = u_-$ , since  $\partial_- u_+ = u_{-+} = u_{+-} = \partial_+ u_-$ . The corresponding spin zero conserved charge is the **topological charge**

$$Q_0 = \int_{-\infty}^{+\infty} dx (u_+ - u_-) = 2 \int_{-\infty}^{+\infty} dx u_x = 2[u]_{-\infty}^{+\infty}.$$

**Note:**  $T_1$  is a total derivative of  $u$ , but  $u$  could have different limits as  $x \rightarrow \pm\infty$ ,<sup>3</sup> in which case the condition (4.15) is not satisfied by  $U = -u$ . Precisely in those cases the topological charge is non-trivial.

$$\boxed{s = 1} \quad T_2 \supset u_{++}, u_+^2,$$

which is a shorthand for:  $T_2$  is a linear combination of  $u_{++}$  and  $u_+^2$ . However  $u_{++} = (u_+)_+$  is a total derivative, and since  $u_+ \rightarrow 0$  as  $x \rightarrow \pm\infty$  we can disregard this term without loss of generality, and consider  $T_2 = u_+^2$ . Then

$$\partial_- T_2 = \partial_- u_+^2 = 2u_+ u_{+-} \underset{\text{EoM}}{=} -2V'(u)u_+ = -2\partial_+ V(u) \equiv \partial_+ X_0$$

with  $X_0 = -2V(u)$ . Therefore

$$\boxed{Q_1 = \int_{-\infty}^{+\infty} dx (T_2 - X_0) = \int_{-\infty}^{+\infty} dx [u_+^2 + 2V(u)]} \quad (4.16)$$

is conserved.

#### **REMARK:**

given a pair of densities  $(T_{s+1}, X_{s-1})$  leading to a conserved charge  $Q_s$ , we can switch the roles of  $x^+$  and  $x^-$  to find a second pair of densities  $(T_{-s-1}, X_{-s+1})$ , which lead to a conserved charge

$$\boxed{Q_{-s} = \int_{-\infty}^{+\infty} dx (T_{-s-1} - X_{-s+1})}$$

of opposite spin.

So at  $|s| = 1$  we have the conserved charges

$$Q_{+1} = \int_{-\infty}^{+\infty} dx [u_+^2 + 2V(u)]$$

$$Q_{-1} = \int_{-\infty}^{+\infty} dx [u_-^2 + 2V(u)].$$

---

<sup>3</sup>The BC's are  $u_t, u_x, V(u), V'(u) \rightarrow 0$  as  $x \rightarrow \pm\infty$ . Depending on the potential,  $u$  could be allowed to have different limits as  $x \rightarrow \pm\infty$ .

Taking the sum and difference and choosing a convenient normalization, we find two conserved charges

$$\begin{aligned} \frac{1}{4}(Q_1 + Q_{-1}) &= \int_{-\infty}^{+\infty} dx \left[ \frac{1}{4}(u_+^2 + u_-^2) + V(u) \right] \\ &\equiv \boxed{E = \int_{-\infty}^{+\infty} dx \left[ \frac{1}{2}u_t^2 + \frac{1}{2}u_x^2 + V(u) \right]} \end{aligned} \quad (4.17)$$

$$\begin{aligned} \frac{1}{4}(Q_{-1} - Q_1) &= \int_{-\infty}^{+\infty} dx \frac{1}{4}(u_-^2 - u_+^2) \\ &\equiv \boxed{P = - \int_{-\infty}^{+\infty} dx u_t u_x}, \end{aligned} \quad (4.18)$$

which are interpreted as the **energy**  $E$  and the **momentum**  $P$ .

$$\boxed{s = 2} \quad T_3 \supset u_{+++}, u_{++}u_+, u_+^3,$$

but  $u_{+++} = (u_{++})_+$  and  $u_{++}u_+ = \frac{1}{2}(u_+^2)_+$  are total derivatives of functions which vanish at spatial infinity, hence they can be disregarded.<sup>4</sup> So without loss of generality we can take  $T_3 = u_+^3$  and then

$$\partial_- T_3 = \partial_- u_+^3 = 3u_+^2 u_{+-} \underset{\text{EoM}}{=} -3V'(u)u_+^2.$$

The RHS of the previous equation **cannot be a total  $x^+$ -derivative**, because the highest  $x^+$  derivative of  $u$  (in this case  $u_+$ ) does not appear linearly.

**\* EXERCISE:** Think about it and convince yourself that this statement is correct. Suppose that  $\partial_+^n u$  is the highest  $x^+$ -derivative of  $u$  appearing in a function  $Y$  of  $u$  and its  $x^+$ -derivatives. How does the highest  $x^+$ -derivative of  $u$  appear in  $\partial_+ Y$  then?

We learn therefore that there is **no conserved charge  $Q_2$  of spin 2** built out of polynomially conserved densities.

$$\boxed{s = 3} \quad T_4 \supset u_{++++}, u_{+++}u_+, u_{++}^2, u_{++}u_+^2, u_+^4,$$

but we can drop the first and fourth term as they are total derivatives of functions which vanish at spatial infinity. Moreover  $u_{+++}u_+ = -u_{++}^2 + (u_{++}u_+)_+$ , so we can also disregard one of  $u_{+++}u_+$  and  $u_{++}^2$  without loss of generality. The most general expression for  $T_4$  up to an irrelevant total  $x^+$ -derivative is therefore

$$\boxed{T_4 = u_{++}^2 + \frac{1}{4}\lambda^2 u_+^4}, \quad (4.19)$$

---

<sup>4</sup>Note that  $u_{+\dots+} \xrightarrow{x \rightarrow \pm\infty} 0$  assuming uniform convergence, which allows us to commute derivatives and limits. (We need to use  $\partial_+ = -\partial_- + 2\partial_+$  and the EoM.)

where  $\lambda$  is a constant to be determined below and the factor of  $1/4$  was inserted for later convenience.<sup>5</sup> Then

$$\begin{aligned}\partial_- T_4 &= 2u_{++}u_{+-} + \lambda^2 u_+^3 u_{+-} \stackrel{\text{EoM}}{=} -2u_{++} (V'(u))_+ - \lambda^2 u_+^3 V'(u) \\ &= -2u_{++}u_+ V''(u) - \lambda^2 u_+^3 V'(u) = -(u_+^2)_+ V''(u) - \lambda^2 u_+^3 V'(u) .\end{aligned}$$

This may not seem very promising, but we can extract a total derivative from the first term using the trick familiar from integration by parts:

$$\begin{aligned}&= -(u_+^2 V''(u))_+ + u_+^3 V'''(u) - \lambda^2 u_+^3 V'(u) \\ &= -(u_+^2 V''(u))_+ + u_+^3 [V'''(u) - \lambda^2 V'(u)] .\end{aligned}\tag{4.20}$$

We are hoping to obtain a total  $x^+$ -derivative. The first term in (4.20) is a total  $x^+$ -derivative, but in the second term the highest derivative  $u_+$  does not appear linearly. By the previous argument which was the topic of the exercise, the second term is a total  $x^+$ -derivative if and only if

$$\boxed{V'''(u) - \lambda^2 V'(u) = 0} .\tag{4.21}$$

If (4.21) holds, we have  $X_2 = -u_+^2 V''(u)$  and

$$\boxed{Q_3 = \int_{-\infty}^{+\infty} dx (T_4 - X_2) = \int_{-\infty}^{+\infty} dx \left[ u_{++}^2 + \frac{1}{4} \lambda^2 u_+^4 + u_+^2 V''(u) \right]}\tag{4.22}$$

is a **conserved charge of spin 3**. If instead (4.16) does not hold, there is no extra (polynomially) conserved charge of spin 3.

To summarize, the **relativistic field theories** which have an **extra conserved charge (of spin 3)**, in addition to the topological charge (if that is non-trivial), the energy and the momentum, are those with a scalar potential  $V(u)$  which satisfies equation (4.21) for some value of the constant  $\lambda$ . Let us examine the various possibilities:

$$1. \quad \boxed{\lambda^2 = 0} : \quad V(u) = A + B(u - u_0)^2,$$

where  $A$  and  $B$  are constants. Up to a linear redefinition of  $u$ , this scalar potential leads to the **Klein-Gordon equation**. This is a linear equation which describes a free field (*i.e.* a field free from interactions) and is therefore not interesting from the point of view of solitons.

$$2. \quad \boxed{\lambda^2 \neq 0} : \quad V(u) = A + B e^{\lambda u} + C e^{-\lambda u},$$

where  $A, B$  and  $C$  are constants.

<sup>5</sup>To be precise,  $T_4$  should be written as a linear combination of  $u_{++}^2$  and  $u_+^4$ . It turns out that the coefficient of  $u_{++}$  must be non-vanishing, hence we can normalise it to 1.

a) If only one of  $B, C$  is non-vanishing, the EoM is either

$$\underline{C = 0}: \quad u_{+-} = -B\lambda e^{\lambda u} \quad \text{or} \quad \underline{B = 0}: \quad u_{+-} = C\lambda e^{-\lambda u}.$$

By a linear redefinition of  $u$ , we can always rewrite the EoM as the **Liouville equation**

$$\boxed{u_{+-} = e^u}. \quad (4.23)$$

b) If neither  $B$  or  $C$  vanish, then by a linear redefinition of  $u$  we can write the EoM as the **sine-Gordon equation**

$$\boxed{u_{+-} = -\sin u} \quad (4.24)$$

if  $\lambda^2 < 0$ , or as the **sinh-Gordon equation**

$$\boxed{u_{+-} = -\sinh u} \quad (4.25)$$

if  $\lambda^2 > 0$ .

The equations (4.23)-(4.25) are **special**: they have “**hidden**” **conservation laws** that generic interacting relativistic field equations  $u_{+-} = -V'(u)$  don't have.

One could carry on and find that these equations have more and more hidden conservation laws. But there are more systematic ways to find them, that you might return to next term. Instead, we will now move on to studying conservations laws for the KdV equation.

## 4.4 Conserved quantities for the KdV equation

Let us return to the KdV equation

$$u_t + 6uu_x + u_{xxx} = 0.$$

We can rewrite the KdV equation as a continuity equation

$$\frac{\partial}{\partial t}u + \frac{\partial}{\partial x}(3u^2 + u_{xx}) = 0$$

and since the BC's appropriate for KdV on the line  $\mathbb{R}$  are that  $u, u_x, u_{xx}, \dots \rightarrow 0$  as  $x \rightarrow \pm\infty$ , we deduce that the **mass** of the water wave<sup>6</sup>

$$\boxed{Q_1 = \int_{-\infty}^{+\infty} dx \, u}, \quad (4.26)$$

---

<sup>6</sup>(4.26) is the (net) area under the profile of the wave, taking  $u = 0$  (flat water surface) as zero. Assuming that water has constant density (mass per unit area) and choosing units so that the density is 1, (4.26) is the mass of the wave.

is **conserved**.

Next, we can ask whether  $\rho = u^2$  is a conserved charge density. Let us compute

$$\begin{aligned}(u^2)_t &= 2uu_t \stackrel{\text{KdV}}{=} -12u^2u_x - 2uu_{xxx} = -4(u^3)_x - 2uu_{xxx} \\ &= (-4u^3 - 2uu_{xx})_x + 2u_xu_{xx} = (-4u^3 - 2uu_x + u_x^2)_x ,\end{aligned}$$

where to go from the first to the second line we used the trick familiar from integration by parts,  $fg_x = (fg)_x - f_xg$ . (We say that  $fg_x$  and  $-f_xg$  are equal up to a total  $x$ -derivative.) Hence we deduce that

$$Q_2 = \int_{-\infty}^{+\infty} dx u^2 , \quad (4.27)$$

which is interpreted as the **momentum** of the wave, is **conserved**.

Next, what about  $\rho = u^3$ ? Using the notation “=” to mean “equal up to a total  $x$ -derivative” and striking out terms which are total derivatives (t.d.), we find

$$\begin{aligned}(u^3)_t &= 3u^2u_t \stackrel{\text{KdV}}{=} -18\cancel{u^3u_x}^{\text{t.d.}} - 3u^2u_{xxx} = ”6uu_xu_{xx}” \\ &\stackrel{\text{KdV}}{=} -u_tu_{xx} - \cancel{u_{xxx}u_{xx}}^{\text{t.d.}} = ”u_{tx}u_x = \frac{1}{2}(u_x^2)_t” ,\end{aligned}$$

so rearranging we find a third **conserved** charge

$$Q_3 = \int_{-\infty}^{+\infty} dx \left( u^3 - \frac{1}{2}u_x^2 \right) , \quad (4.28)$$

which is interpreted as the **energy** of the wave.

It turns out that the conservation laws (4.26)-(4.28) of **mass**, **momentum** and **energy** follow by Noether’s theorem from the “obvious” symmetries

$$\begin{array}{lll} u \mapsto u + c & \implies & \text{mass conservation} \\ x \mapsto x + c' & \implies & \text{momentum conservation} \\ t \mapsto t + c'' & \implies & \text{energy conservation} \end{array}$$

of the KdV equation, so they are expected. But then surprisingly [Miura et al., 1968] found (by hand!) **eight more conserved charges**, all (but one, see [Ex 23]) of the form

$$Q_n = \int_{-\infty}^{+\infty} dx (u^n + \dots) ,$$

e.g.

$$\begin{aligned}
 Q_4 &= \int_{-\infty}^{+\infty} dx \left( u^4 - 2uu_x^2 + \frac{1}{5}u_{xx}^2 \right) \\
 Q_5 &= \int_{-\infty}^{+\infty} dx \left( u^5 - 5u^2u_x^2 + uu_{xx}^2 - \frac{1}{14}u_{xxx}^2 \right) \\
 &\vdots \\
 Q_{10} &= \int_{-\infty}^{+\infty} dx \left( u^{10} - 60u^7u_x^2 + (29 \text{ terms}) + \frac{1}{4862}u_{xxxxxxx}^2 \right).
 \end{aligned} \tag{4.29}$$

**\* EXERCISE:** Calculate  $Q_6, \dots, Q_9$  as well and the 29 missing terms in  $Q_{10}$ .

7

This surprising result raises **two natural questions**:

1. Are there infinitely many more conserved charges?
2. If so, is there a systematic way to find them?

## 4.5 The Gardner transform

The answer to both questions is affirmative, and is based on a very clever (though at first sight unintuitive) method devised by **Gardner** [Miura et al., 1968].

First, let us suppose that the KdV field  $u(x, t)$  can be expressed in terms of another function  $v(x, t)$  as

$$u = \lambda - v^2 - v_x, \tag{4.30}$$

where  $\lambda$  is a real parameter. Substituting (4.30) in the KdV equation we find

$$\begin{aligned}
 0 &= (\lambda - v^2 - v_x)_t + 6(\lambda - v^2 - v_x)(\lambda - v^2 - v_x)_x + (\lambda - v^2 - v_x)_{xxx} \\
 &= \dots \quad \text{[Ex 24]} \\
 &= - \left( 2v + \frac{\partial}{\partial x} \right) [v_t + 6(\lambda - v^2)v_x + v_{xxx}] = 0.
 \end{aligned} \tag{4.31}$$

So

$$\boxed{\text{KdV for } u \iff (4.31) \text{ for } v},$$

---

<sup>7</sup>Just kidding.

and in particular, if  $v$  solves

$$\boxed{v_t + 6(\lambda - v^2)v_x + v_{xxx} = 0} , \quad (4.32)$$

then  $u$  given by (4.30) solves KdV.

For  $\lambda = 0$ , (4.32) is the “wrong sign” mKdV equation that you encountered in **[Ex 13.2]**, and

$$\boxed{u = -v^2 - v_x} \quad (4.33)$$

was known as the **Miura transform**, found by Miura earlier in 1968 [Miura, 1968].

**Gardner’s idea** was to bring the equation for  $v$  closer to the KdV equation when  $\lambda \neq 0$ . This is achieved by setting

$$\boxed{\begin{aligned} v &= \epsilon w + \frac{1}{2\epsilon} \\ \lambda &= \frac{1}{4\epsilon^2} \end{aligned}} \quad (4.34)$$

for some non-vanishing real constant  $\epsilon$ . Then

$$\lambda - v^2 = \frac{1}{4\epsilon^2} - \left( \epsilon w + \frac{1}{2\epsilon} \right)^2 = -w - \epsilon^2 w^2 ,$$

which implies that  $u$  and  $w$  are related by the **Gardner transform (GT)**

$$\boxed{u = -w - \epsilon w_x - \epsilon^2 w^2} . \quad (4.35)$$

We will use the free parameter  $\epsilon$  to great advantage below.

In terms of  $w$ , the **KdV equation for  $u$** , or equivalently equation (4.31) for  $v$  becomes

$$\left( 2\epsilon w + \frac{1}{\epsilon} + \frac{\partial}{\partial x} \right) [\epsilon w_t - 6(w + \epsilon^2 w^2)\epsilon w_x + \epsilon w_{xxx}] = 0 ,$$

or equivalently

$$\boxed{\left( 1 + \epsilon \frac{\partial}{\partial x} + 2\epsilon^2 w \right) [w_t - 6(w + \epsilon^2 w^2)w_x + w_{xxx}] = 0} . \quad (4.36)$$

In particular, any  $w$  that solves the simpler equation

$$\boxed{w_t - 6(w + \epsilon^2 w^2)w_x + w_{xxx} = 0} \quad (4.37)$$

produces a  $u$  that solves the KdV equation by the Gardner transform (4.35).

Now we are going to think about this backwards: let’s view  $u$  as a **fixed solution of KdV**, while  $w$  **varies with**  $\epsilon$  so that (4.35) holds. Then



- For  $\epsilon = 0$ , equation (4.36) is nothing but the KdV equation with reversed middle terms. Indeed the Gardner transform reduces to  $u = -w$  in this case.<sup>8</sup> (Incidentally, this shows that we can extend the domain of  $\epsilon$  to include 0.)
- For  $\epsilon \neq 0$ , we encounter **two problems**:

1. To obtain  $w$  in terms of  $u$ , we need to solve a **differential equation** (4.35);
2. The **differential operator**  $1 + \epsilon \frac{\partial}{\partial x} + 2\epsilon^2 w$  in (4.36) is **non-trivial**. It might have a non-vanishing kernel, so we can't immediately conclude that (4.37) holds.

Gardner's intuition was that we can **solve both problems at once** by viewing  $w$  as a **formal power series in  $\epsilon$** :<sup>9</sup>

$$w(x, t) = \sum_{n=0}^{\infty} w_n(x, t) \epsilon^n = w_0(x, t) + w_1(x, t) \epsilon + w_2(x, t) \epsilon^2 + \dots \quad (4.38)$$

1. To solve the first problem, we substitute (4.38) in the Gardner transform (4.35)

$$\begin{aligned} u &= -(w_0 + w_1 \epsilon + w_2 \epsilon^2 + \dots) - \epsilon(w_0 + w_1 \epsilon + w_2 \epsilon^2 + \dots)_x \\ &\quad - \epsilon^2(w_0 + w_1 \epsilon + w_2 \epsilon^2 + \dots)^2 \\ &= -w_0 \quad -\epsilon w_1 \quad -\epsilon^2 w_2 \quad -\epsilon^3 w_3 \quad + \dots \\ &\quad \quad -\epsilon w_{0,x} \quad -\epsilon^2 w_{1,x} \quad -\epsilon^3 w_{2,x} \quad + \dots \\ &\quad \quad \quad -\epsilon^2 w_0^2 \quad -\epsilon^3 2w_0 w_1 \quad + \dots \end{aligned}$$

and invert it to determine  $w$  in terms of  $u$ . Since  $u$  is fixed, it is of order  $\epsilon^0$ . Comparing order by order we obtain:

$$\begin{aligned} \epsilon^0 : & \quad w_0 = -u \\ \epsilon^1 : & \quad w_1 = -w_{0,x} = u_x \\ \epsilon^2 : & \quad w_2 = -w_{1,x} - w_0^2 = -u_{xx} - u^2 \\ \epsilon^3 : & \quad w_3 = -w_{2,x} - 2w_0 w_1 = u_{xxx} + 4u u_x \\ & \quad \vdots \end{aligned} \quad (4.39)$$

which in principle determines recursively all the coefficients  $w_n$  of the formal power series (4.38) in terms of  $u$ .

<sup>8</sup>Sorry, I should have taken  $w \rightarrow -w$  in (4.34). Too late to change that now...

<sup>9</sup>By a formal power series we mean that we don't worry about the convergence of the series. (4.38) is actually an asymptotic expansion, for those who know what that is.

2. Since  $w$  is a formal power series in  $\epsilon$ , so is the expression inside the square brackets in (4.36):

$$[w_t - 6(w + \epsilon^2 w^2)w_x + w_{xxx}] \equiv z(x, t) = \sum_{n=0}^{\infty} z_n(x, t) \epsilon^n = z_0 + z_1 \epsilon + z_2 \epsilon^2 + \dots$$

The same applies to the differential operator

$$A \equiv \mathbb{1} + \epsilon \frac{\partial}{\partial x} + 2\epsilon^2 w \equiv \mathbb{1} + \sum_{n=1}^{\infty} A_n \epsilon^n,$$

where  $\mathbb{1}$  is the identity operator, and  $A_n$  are linear (differential) operators:

$$A_1 = \frac{\partial}{\partial x}, \quad A_2 = 2w_0 \cdot, \quad A_3 = 2w_1 \cdot, \quad A_4 = 2w_2 \cdot, \quad \dots$$

where I wrote the dots to make clear which operators act by multiplication by a function. Then (4.36) becomes the formal power series equation

$$\begin{aligned} 0 &= \left(1 + \sum_{n=1}^{\infty} A_n \epsilon^n\right) \left(\sum_{k=0}^{\infty} z_k \epsilon^k\right) \\ &= z_0 \quad + \epsilon z_1 \quad + \epsilon^2 z_2 \quad + \epsilon^3 z_3 \quad + \dots \\ &\quad + \epsilon A_1 z_0 \quad + \epsilon^2 A_1 z_1 \quad + \epsilon^3 A_1 z_2 \quad + \dots \\ &\quad \quad + \epsilon^2 A_2 z_0 \quad + \epsilon^3 A_2 z_1 \quad + \dots \\ &\quad \quad \quad + \epsilon^3 A_3 z_0 \quad + \dots \\ &\quad \quad \quad \quad + \dots \end{aligned}$$

which we can solve order by order as follows:

$$\begin{aligned} \epsilon^0 : & \quad z_0 = 0 \\ \epsilon^1 : & \quad z_1 = -A_1 z_0 \\ \epsilon^2 : & \quad z_2 = -A_1 z_1 - A_2 z_0 = 0 \\ \epsilon^3 : & \quad z_3 = -A_1 z_2 - A_2 z_1 - A_3 z_0 = 0 \\ & \quad \vdots \end{aligned} \tag{4.40}$$

We found that, **order by order** in the formal power series in  $\epsilon$ , **equation (4.37) holds!** But (4.37) is a continuity equation

$$\boxed{\frac{\partial}{\partial t} w + \frac{\partial}{\partial x} (-3w^2 - 2\epsilon^2 w^3 + w_{xx}) = 0} . \tag{4.41}$$

Since  $w, w_x, w_{xx}, \dots \rightarrow 0$  as  $x \rightarrow \pm\infty$  order by order in powers of  $\epsilon$ , the charge

$$\boxed{\tilde{Q} = \int_{-\infty}^{+\infty} dx \, w} \tag{4.42}$$

is **conserved**.

Now comes the important point: since  $w = \sum_{n=0}^{\infty} w_n \epsilon^n$  is a formal power series in  $\epsilon$ , so is the conserved charge  $\tilde{Q}$  too:<sup>10</sup>

$$\tilde{Q} = \int_{-\infty}^{+\infty} dx \sum_{n=0}^{\infty} w_n \epsilon^n = \sum_{n=0}^{\infty} \epsilon^n \int_{-\infty}^{+\infty} dx w_n \equiv \sum_{n=0}^{\infty} \epsilon^n \tilde{Q}_n .$$

And since  $\tilde{Q}$  is a conserved charge for all values of the free parameter  $\epsilon$ , it must be that the **charges**

$$\boxed{\tilde{Q}_n = \int_{-\infty}^{+\infty} dx w_n} \quad (n = 0, 1, 2, \dots) \quad (4.43)$$

are **all separately conserved!**

Going back to (4.39), we find that the first few conserved charges are

$$\boxed{\begin{aligned} \tilde{Q}_0 &= - \int_{-\infty}^{+\infty} dx u \equiv -Q_1 \\ \tilde{Q}_1 &= + \int_{-\infty}^{+\infty} dx u_x = [u]_{-\infty}^{+\infty} = 0 \\ \tilde{Q}_2 &= - \int_{-\infty}^{+\infty} dx (u_{xx} + u^2) = - \int_{-\infty}^{+\infty} dx u^2 \equiv -Q_2 \\ \tilde{Q}_3 &= + \int_{-\infty}^{+\infty} dx (u_{xxx} + 4uu_x) = [u_{xx} + 2u^2]_{-\infty}^{+\infty} = 0 \\ &\vdots \end{aligned}} \quad (4.44)$$

As you might have guessed, the **general pattern** is as follows:

$$\begin{aligned} \tilde{Q}_{2n-1} &= \int_{-\infty}^{+\infty} dx (\text{total derivative}) = 0 \\ \tilde{Q}_{2n-2} &= \text{const} \times Q_n = \text{const} \times \int_{-\infty}^{+\infty} dx (u_n + \dots) \neq 0 . \end{aligned}$$

See [Drazin and Johnson, 1989] for a general proof.

The existence of **infinitely many conserved charges** makes the **KdV** equation “**integrable**”. As you’ll start seeing in the exercises for this chapter, these unexpected conservation laws give us a lot of information about multi-soliton solutions of the KdV equation, see **[Ex 23]** and **[Ex 25]**.

---

<sup>10</sup>Strictly speaking the middle equality assumes convergence, but we are working with a formal expansion, so we don’t need to worry about this subtlety.

# Chapter 5

## The Bäcklund transform

The main reference for this chapter is §5.4 of [Drazin and Johnson, 1989].

**So far**, we have constructed **exact** analytic solutions for moving solitons only as **travelling waves**, which describe the propagation of a **single soliton**. Our **next** goal will be to construct **exact** analytic solutions for **multiple colliding solitons**. The method that we will use in this chapter is a solution-generating technique called the **Bäcklund transform**.

The method was introduced in the late 19th century by the Swedish mathematician **Albert Victor Bäcklund** and by the Italian mathematician **Luigi Bianchi**<sup>1</sup> in the 1880s to map a pair of surfaces tangent to one another in three-dimensional space into another pair of surfaces tangent to one another in a second three-dimensional space. The sine-Gordon equation happens to appear in this context when one considers hyperboloids, which are surfaces of negative curvature.

There are **two main uses of the Bäcklund transform**:

1. To generate **solutions of a difficult PDE from solutions of a simpler PDE**;
2. To generate **new solutions of a given PDE from already known solutions of the same PDE**.

We will mostly be interested in use 2, but you will see examples of use 1 in Ex 26-28 in the

---

<sup>1</sup>who, notably, was born in my hometown, Parma. Please remember this for the exam ;) This is the same Bianchi after whom the Bianchi identities in differential geometry and general relativity are named.

problem sheet. Our final goal in this chapter will be to obtain multi-soliton solutions of the sine-Gordon equation.

## 5.1 Definition

Consider two functions  $u$  and  $v$ , and two differential equations

$$\boxed{P[u] = 0} \quad (5.1)$$

$$\boxed{Q[v] = 0} \quad (5.2)$$

where  $P$  and  $Q$  are two differential operators.

If there is a pair of relations (which could be differential equations)

$$\boxed{R_1[u, v] = 0, \quad R_2[u, v] = 0} \quad (5.3)$$

between  $u$  and  $v$  such that

- If  $P[u] = 0$ , then (5.3) can be solved for  $v$ , to give a solution of  $Q[v] = 0$ ;
- If  $Q[v] = 0$ , then (5.3) can be solved for  $u$ , to give a solution of  $P[u] = 0$ ;

then (5.3) is called a **Bäcklund transform (BT)**. If furthermore  $P = Q$ , so that the two differential equations are identical, then (5.3) is called an **auto-Bäcklund transform (a-BT)**.

This is useful if (5.3) is easier to solve than (5.1) or (5.2). Then we can use (5.3) to generate solutions of the harder equation from solutions of the easier equation. If  $P = Q$ , we can start from a simple seed solution (e.g.  $u = 0$ ) to generate new non-trivial solutions.

### Vocabulary:

(5.1) and (5.2) are “**integrability conditions**” for the Bäcklund transform relations (5.3).

(5.3) can be integrated for  $v$  if the integrability condition  $P[u] = 0$  is satisfied.

(5.3) can be integrated for  $u$  if the integrability condition  $Q[v] = 0$  is satisfied.

## 5.2 A simple example

Take the two-dimensional Laplace operator  $P = Q = \partial_x^2 + \partial_y^2$  in (5.1) and (5.2):

$$P[u] = u_{xx} + u_{yy} = 0 \quad (5.4)$$

$$Q[v] = v_{xx} + v_{yy} = 0 \quad (5.5)$$

and for the Bäcklund transform (5.3)

$$\begin{aligned} R_1[u, v] &= u_x - v_y = 0 \\ R_2[u, v] &= u_y + v_x = 0. \end{aligned} \tag{5.6}$$

Let us check that (5.4)-(5.5) are integrability conditions for (5.6). Differentiating (5.6) we find

$$\begin{aligned} 0 &= +\partial_x R_1 + \partial_y R_2 = +u_{xx} - v_{yx} + u_{yy} + v_{xy} = u_{xx} + u_{yy} \\ 0 &= -\partial_y R_1 + \partial_x R_2 = -u_{xy} + v_{yy} + u_{yx} + v_{xx} = v_{xx} + v_{yy}, \end{aligned}$$

therefore the relations (5.6) imply (5.4) and (5.5).<sup>2</sup> This shows that (5.6) is an **auto-Bäcklund transform** for the two-dimensional Laplace equation.

### EXAMPLE:

$v(x, y) = 2xy$  solves the Laplace equation (5.5). Let us use the a-BT to find another solution  $u$  of the same equation:

$$\begin{cases} u_x = v_y = 2x \\ u_y = -v_x = -2y \end{cases} \implies \begin{cases} u = x^2 + f(y) \\ f'(y) = -2y \end{cases} \implies f(y) = -y^2 + c,$$

so we find the function  $u(x, y) = x^2 - y^2 + c$ , where  $c$  is a constant. It is immediate to check that this  $u$  solves the Laplace equation (5.4).

The equations  $R_1[u, v] = R_2[u, v] = 0$  in (5.6) are nothing but the **Cauchy-Riemann equations** for the **holomorphic** (= complex analytic) **function**  $w = u + iv$  of the complex variable  $z = x + iy$ . In the example above,  $w(z) = z^2 + c$ . The equations  $P[u] = 0$  and  $Q[v] = 0$  in (5.4)-(5.5) simply state that the real and imaginary parts of a holomorphic function are harmonic, that is, they solve the Laplace equation. Two such functions  $u$  and  $v$  are often called **harmonic conjugate** of each other.

### REMARKS:

1. **Given**  $v$ , the Bäcklund transform (5.6) is a system of **two equations for**  $u$ . Generically there wouldn't be any solutions for the system (5.6). For example, if we pick  $v = x^2$ , then the system

$$\begin{cases} u_x = v_y = 0 \\ u_y = -v_x = -2x \end{cases}$$

has no solutions for  $u$ . But  $v = x^2$  doesn't solve (5.5)! The **integrability condition** (5.5) is what guarantees that the system (5.6) can be consistently solved for  $u$ .

---

<sup>2</sup>Note: in this example we don't even need to use the other differential equation. This is not always the case.

2. This **auto-Bäcklund transform** generates a new solution to the Laplace equation from a seed solution, but if we apply it a second time we get back the original seed solution (up to an irrelevant integration constant that we can ignore). So this auto-Bäcklund transform is an **involution**. To get further solutions we will need to introduce a parameter.

### 5.3 Bäcklund transform for sine-Gordon

Recall that the sine-Gordon equation written in light-cone coordinates  $x^\pm = \frac{1}{2}(t \pm x)$  is

$$\boxed{u_{+-} = -\sin u} . \quad (5.7)$$

Let us try the Bäcklund transform

$$\boxed{\begin{aligned} (u-v)_+ &= \frac{2}{a} \sin\left(\frac{u+v}{2}\right) \\ (u+v)_- &= -2a \sin\left(\frac{u-v}{2}\right) \end{aligned}} \quad (5.8)$$

with the **parameter**  $a$ . Cross-differentiating,<sup>3</sup>

$$\begin{aligned} (u-v)_{+-} &= \frac{1}{a} \cos\left(\frac{u+v}{2}\right) \cdot (u+v)_- = -2 \cos\left(\frac{u+v}{2}\right) \sin\left(\frac{u-v}{2}\right) \\ &= -\sin u + \sin v \\ (u+v)_{-+} &= -a \cos\left(\frac{u-v}{2}\right) \cdot (u-v)_+ = -2 \cos\left(\frac{u-v}{2}\right) \sin\left(\frac{u+v}{2}\right) \\ &= -\sin u - \sin v . \end{aligned}$$

Adding and subtracting, we find that both  $u$  and  $v$  obey the sine-Gordon equation:

$$\boxed{u_{+-} = -\sin u} \quad (5.9)$$

$$\boxed{v_{+-} = -\sin v} \quad (5.10)$$

Therefore (5.8) is an **auto-Bäcklund transform for the sine-Gordon equation, for any value of  $a$** . The extra parameter will allow us to generate multi-soliton solutions of the sine-Gordon equation. We will start by rederiving the one-kink solution using the auto-Bäcklund transform (5.8).

---

<sup>3</sup>Recall:  $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$ , which implies  $\sin(A+B) + \sin(A-B) = 2 \sin A \cos B$ .

## 5.4 First example: the sine-Gordon kink from the vacuum

Let us take the **vacuum solution**

$$\boxed{v = 0} \quad (5.11)$$

as our **seed solution**. Then the auto-Bäcklund transform (5.8) is

$$\begin{aligned} u_+ &= \frac{2}{a} \sin \frac{u}{2} \\ u_- &= -2a \sin \frac{u}{2}. \end{aligned} \quad (5.12)$$

We can integrate both equations by separation of variables, using the indefinite integral

$$\int \frac{du}{\sin \frac{u}{2}} = 2 \log \tan \frac{u}{4}$$

up to an integration constant. We get

$$\begin{cases} \frac{2}{a}x^+ = 2 \log \tan \frac{u}{4} + f(x^-) \\ -2ax^- = 2 \log \tan \frac{u}{4} + g(x^+) \end{cases} \quad (5.13)$$

where the functions  $f$  and  $g$  are “constants” of integration. They are only constant with respect to the variable that is integrated, but they can (and do!) depend on the other variable.

Subtracting and rearranging, we get

$$\frac{2}{a}x^+ + g(x^+) = -2ax^- + f(x^-). \quad (5.14)$$

The left-hand-side is only a function of  $x^+$ . The right-hand-side is only a function of  $x^-$ . But the two sides are equal, so they must be equal to a constant, that we set to be  $-2c$  for future convenience. Therefore

$$\begin{aligned} f(x^-) &= 2ax^- - 2c \\ g(x^+) &= -\frac{2}{a}x^+ - 2c \end{aligned}$$

and so

$$2 \log \tan \frac{u}{4} = \frac{2}{a}x^+ - 2ax^- + 2c,$$

that is

$$\boxed{u = 4 \arctan \left( e^{\frac{1}{a}x^+ - ax^- + c} \right)}. \quad (5.15)$$

Finally, we convert to  $(x, t)$  coordinates:

$$\frac{1}{a}x^+ - ax^- = \frac{1}{2a}(t+x) - \frac{a}{2}(t-x) = \frac{1}{2} \left[ \left(a + \frac{1}{a}\right)x - \left(a - \frac{1}{a}\right)t \right] = \frac{1+a^2}{2a} \left( x - \frac{a^2+1}{a^2-1}t \right).$$



Defining

$$\left. \begin{aligned} v &:= \frac{a^2 - 1}{a^2 + 1} \\ \epsilon &:= \text{sign}(a) \\ \gamma &:= \frac{1}{\sqrt{1 - v^2}} \stackrel{*Ex}{=} \frac{1 + a^2}{2|a|} \end{aligned} \right\}, \quad (5.16)$$





the solution (5.15) generated by auto-Bäcklund transform of the vacuum is

$$u(x, t) = 4 \arctan \left( e^{\epsilon \gamma (x - x_0 - vt)} \right), \quad (5.17)$$

where we traded the integration constant  $c$  for  $x_0$ . This solution describes a kink or an anti-kink moving at velocity  $v$ .

**Properties:**


$$\begin{array}{ll} a > 0: & \text{kink} \\ a < 0: & \text{anti-kink} \end{array} \quad \begin{array}{ll} |a| > 1: & \text{right-moving} \\ |a| < 1: & \text{left-moving} \end{array}$$

$a < -1$ :	$-1 < a < 0$	$0 < a < 1$	$a > 1$
Right-moving anti-kink	Left-moving anti-kink	Left-moving kink	Right-moving kink
			

So the auto-Bäcklund transform creates a kink/anti-kink from the vacuum! By varying the parameter  $a \in \mathbb{R} \setminus \{0\}$  and the integration constant  $x_0$  or  $c$ , we reproduce all the kink and anti-kink solutions derived in section 2.2 as travelling waves.


The **amazing fact** is that this holds more generally: the auto-Bäcklund transform always adds a kink or an anti-kink to the seed solution.<sup>4</sup> (The only exception is if one tries to add a soliton with the same velocity as one already present.) Therefore we can think of the **auto-Bäcklund transform** as a **solution-generating technique which “adds” kinks or anti-kinks**.

We will use the following graph to denote the action of a Bäcklund transform on with parameter  $a$  and integration constant  $c$  on a seed solution  $u_1$ , which adds a kink or anti-kink and generates the new solution  $u_2$ :



<sup>4</sup>Which of the two is added depends on the seed. More about this later.

We can add a kink/anti-kink wherever we like (by choosing  $c$ ) and with whatever velocity we like (by choosing  $a$ ). For example




adds three kinks/anti-kinks to the seed solution  $u_0$ .

The problem with this is that the **integrations get harder and harder** as we keep “adding” solitons. Luckily, a nice theorem tells us that, having found one-soliton solutions, we can obtain multi-soliton solutions without doing any further integrals.

## 5.5 The theorem of permutability

Let’s **apply the Bäcklund transform twice**, with parameters  $a_1$  and  $a_2$ , in the two possible orders:




The final results  $u_3$  and  $u_4$  of the double Bäcklund transforms both look like the seed solution  $u_0$  with two added solitons, with parameters  $a_1$  and  $a_2$ . Could they actually be the same solution? The answer is yes, according to the following theorem:

**THEOREM** (Bianchi 1902):

For any  $u_1$  and  $u_2$ , the integration constants in the second Bäcklund transforms, which generate  $u_3$  and  $u_4$ , can be arranged such that  $u_3$  and  $u_4$  are equal.


In other words, the  $a_1$  and  $a_2$  BT’s can be made to **commute**. Diagrammatically:



I will spare you the proof of the theorem, which is a bit involved. Hopefully the statement makes intuitive sense, given the soliton content of  $u_3$  and  $u_4$ .

This result has a **nice application**. We have **two ways of getting to  $u_3$  from  $u_0$** : either through  $u_1$  or through  $u_2$ . By comparing these two ways we will be able to get rid of all derivatives in the Bäcklund transform and hence to obtain an **algebraic relation** between the four solutions  $u_0, u_1, u_2, u_3$ .

Let's start by considering the  $\partial_+$  part of the Bäcklund transform, and let's look at the upper route in the previous diagram first:




We have

$$\begin{aligned} (u_1 - u_0)_+ &= \frac{2}{a_1} \sin \frac{u_1 + u_0}{2} \\ (u_3 - u_1)_+ &= \frac{2}{a_2} \sin \frac{u_3 + u_1}{2} . \end{aligned} \tag{5.18}$$

Adding the two equations to cancel  $u_1$  out in the left-hand side, we get

$$\boxed{(u_3 - u_0)_+ = \frac{2}{a_1} \sin \frac{u_1 + u_0}{2} + \frac{2}{a_2} \sin \frac{u_3 + u_1}{2}} . \tag{5.19}$$

For the lower route



we swap  $a_1 \leftrightarrow a_2$ ,  $u_1 \leftrightarrow u_2$  and get


$$\boxed{(u_3 - u_0)_+ = \frac{2}{a_2} \sin \frac{u_2 + u_0}{2} + \frac{2}{a_1} \sin \frac{u_3 + u_2}{2}}. \quad (5.20)$$

We have found two different expressions for  $(u_3 - u_0)_+$ . Equating them, we obtain an **algebraic relation** between  $u_0, u_1, u_2, u_3$ :

$$\boxed{\frac{1}{a_1} \sin \frac{u_1 + u_0}{2} + \frac{1}{a_2} \sin \frac{u_3 + u_1}{2} = \frac{1}{a_2} \sin \frac{u_2 + u_0}{2} + \frac{1}{a_1} \sin \frac{u_3 + u_2}{2}}. \quad (5.21)$$

This is very useful: starting from  $u_0$  equal to the vacuum and two one-soliton solutions  $u_1, u_2$ , we can generate a 2-soliton solution  $u_3$  algebraically. We can then iterate the procedure and get a 3-soliton solution, then a 4-soliton solution, and so on and so forth. What we have found is akin to a “**non-linear superposition principle**”: the Bäcklund transform and the permutability theorem provide us with a machinery to “add” solutions of a non-linear equation!

To **check** that the previous procedure is consistent, let's see what happens for the  $\partial_-$  part of the Bäcklund transform. For the upper route




we have

$$\begin{aligned} (u_1 + u_0)_- &= -2a_1 \sin \frac{u_1 - u_0}{2} \\ (u_3 + u_1)_- &= -2a_2 \sin \frac{u_3 - u_1}{2}. \end{aligned} \quad (5.22)$$

Subtracting the two equations we get

$$\boxed{(u_0 - u_3)_- = 2a_2 \sin \frac{u_3 - u_1}{2} - 2a_1 \sin \frac{u_1 - u_0}{2}}. \quad (5.23)$$

For the lower route



we swap again  $a_1 \leftrightarrow a_2, u_1 \leftrightarrow u_2$  and get

$$(u_0 - u_3)_- = 2a_1 \sin \frac{u_3 - u_2}{2} - 2a_2 \sin \frac{u_2 - u_0}{2} . \quad (5.24)$$

Equating (5.23) and (5.24), we find the algebraic relation

$$a_2 \sin \frac{u_3 - u_1}{2} - a_1 \sin \frac{u_1 - u_0}{2} = a_1 \sin \frac{u_3 - u_2}{2} - a_2 \sin \frac{u_2 - u_0}{2} . \quad (5.25)$$

Consistency requires that the two algebraic relations (5.21) and (5.25) agree. To see that, let's first massage (5.21) into the following form:

$$\frac{1}{a_1} \left( \sin \frac{u_1 + u_0}{2} - \sin \frac{u_3 + u_2}{2} \right) = \frac{1}{a_2} \left( \sin \frac{u_2 + u_0}{2} - \sin \frac{u_3 + u_1}{2} \right) .$$

Multiplying by  $a_1 a_2 / 2$  and using the trigonometric identity  $\sin A \pm \sin B = 2 \sin \frac{A \pm B}{2} \cos \frac{A \mp B}{2}$ , this becomes

$$\boxed{a_2 \sin \frac{u_1 + u_0 - u_3 - u_2}{4} \cos \frac{u_1 + u_0 + u_3 + u_2}{4} = a_1 \sin \frac{u_2 + u_0 - u_3 - u_1}{4} \cos \frac{u_2 + u_0 + u_3 + u_1}{4}} \quad (5.26)$$

where we are allowed to simplify the common cosine factor in the two sides because the argument is a function of  $x$  and  $t$  which is generically different from  $\pi/2$  modulo  $2\pi$ .

Similarly, (5.25) can be massaged to

$$a_1 \left( \sin \frac{u_3 - u_2}{2} + \sin \frac{u_1 - u_0}{2} \right) = a_2 \left( \sin \frac{u_3 - u_1}{2} + \sin \frac{u_2 - u_0}{2} \right) ,$$

which upon using the same trigonometric identity as above becomes

$$\boxed{a_1 \sin \frac{u_3 - u_2 + u_1 - u_0}{4} \cos \frac{u_3 - u_2 - u_1 + u_0}{4} = a_2 \sin \frac{u_3 - u_1 + u_2 - u_0}{4} \cos \frac{u_3 - u_1 - u_2 + u_0}{4}} \quad (5.27)$$

which agrees with equation (5.26) upon simplification. So everything is consistent.

To conclude this discussion, let's massage (the simplified version of) equation (5.26) a bit further, with the aim of determining  $u_3$  given  $u_0, u_1$  and  $u_2$ . Letting  $A = (u_0 - u_3)/4$  and  $B = (u_1 - u_2)/4$ , (5.26) becomes

$$\begin{aligned} a_1 \sin(A - B) &= a_2 \sin(A + B) \\ \implies a_1(\sin A \cos B - \sin B \cos A) &= a_2(\sin A \cos B + \sin B \cos A) . \end{aligned}$$

Dividing through by  $\cos A \cos B$ , we find

$$\begin{aligned} a_1(\tan A - \tan B) &= a_2(\tan A + \tan B) . \\ \implies (a_1 - a_2) \tan A &= (a_1 + a_2) \tan B . \end{aligned}$$

In terms of  $u_0, u_1, u_2, u_3$ , this reads

$$\boxed{\tan \frac{u_0 - u_3}{4} = \frac{a_1 + a_2}{a_1 - a_2} \tan \frac{u_1 - u_2}{4}} , \quad (5.28)$$

which is an improvement on (5.26) since  $u_3$  appears only once. Equivalently, we can write

$$\boxed{\tan \frac{u_3 - u_0}{4} = \frac{a_2 + a_1}{a_2 - a_1} \tan \frac{u_1 - u_2}{4}} . \quad (5.29)$$

Either of (5.28) or (5.29) allow us to express  $u_3$  in terms of  $u_0, u_1, u_2$ .

## 5.6 The two-soliton solution

This is the first nice application of the permutability theorem. Take the vacuum as the seed solution, *i.e.*  $u_0 = 0$ . Then  $u_1$  and  $u_2$  are 1-soliton (*i.e.* kink or antikink) solutions

$$\boxed{\tan \frac{u_i}{4} = e^{\theta_i}} \quad (i = 1, 2) \quad (5.30)$$

where

$$\boxed{\theta_i = \frac{x^+}{a_i} - a_i x^- + c_i = \epsilon_i \gamma_i (x - \bar{x}_i - v_i t)} , \quad (5.31)$$

as seen in section 5.4. Here  $\bar{x}_{1,2}$  are the centres of the two solitons at  $t = 0$ .

Then equation (5.29) gives the double Bäcklund transform  $u_3$ :

$$\tan \frac{u_3}{4} = \mu \tan \frac{u_1 - u_2}{4} = \mu \frac{\tan \frac{u_1}{4} - \tan \frac{u_2}{4}}{1 + \tan \frac{u_1}{4} \tan \frac{u_2}{4}} , \quad (5.32)$$

where

$$\mu = \frac{a_2 + a_1}{a_2 - a_1} \quad (5.33)$$

and we used the trigonometric identity

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \cdot \tan B}$$

in the second equality in equation (5.32). Substituting equation (5.30) in equation (5.32) we find the **2-SOLITON SOLUTION**

$$\tan \frac{u_3}{4} = \mu \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}}. \quad (5.34)$$

**REMARK:**

If the two solitons have the **same velocity**  $v_1 = v_2$ , which means

$$\frac{a_1^2 - 1}{a_1^2 + 1} = \frac{a_2^2 - 1}{a_2^2 + 1} \quad \implies \quad a_1 = \pm a_2 ,$$


then  $\mu = 0, \infty$  and the **2-soliton solution** (5.34) **breaks down**. In particular, there is **no static 2-soliton solution!** As we will see later, this is because the two solitons exert a force on one another.

But this is too fast. We haven't confirmed yet that equation (5.34) contains two solitons. Let's understand that next.

## 5.7 Asymptotics of multisoliton solutions

We will focus here on the 2-soliton solution of the sine-Gordon equation, but the method applies more generally to any multi-soliton solutions of integrable equations (e.g. the KdV equation).

Our **goal** will be to study the new solution (5.34) and **identify two solutions hidden in its asymptotics** for  $t \rightarrow \mp\infty$ , namely BEFORE and AFTER the collision. Here is an example of what the solution may look like at early times (before the collision) and at late times (after the collision) in the case of a collision of a kink and an anti-kink:



It is not completely obvious how to find the early time and late time asymptotics analytically. If we just take  $t \pm \infty$  with  $x$  fixed, the two solitons will be at spatial infinity and we will miss them (unless one of the two has zero velocity, in which case we will see that soliton). We should instead follow one or the other soliton by letting

$$\boxed{t \rightarrow \pm\infty \quad \text{with} \quad X_V = x - Vt \quad \text{fixed}}, \quad (5.35)$$

for some appropriate constant velocity  $V$ . If there is a soliton moving at velocity  $V$  in the original  $(x, t)$  coordinates, it will appear stationary in the  $(X_V, t)$  coordinates. For this reason  $(X_V, t)$  is called a “**comoving frame**”: they are coordinates for a reference frame which moves together with an object (e.g. a soliton) of velocity  $V$ .

Let us try this for the solution (5.34) which we obtained from a double Bäcklund transform of the vacuum. We will now use  $u$  to denote the field in the resulting solution, which reads

$$\tan \frac{u}{4} = \mu \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}}$$

with

$$\mu = \frac{a_2 + a_1}{a_2 - a_1}, \quad \theta_i = \epsilon_i \gamma_i (x - v_i t - \bar{x}_i).$$

If we switch to a comoving frame with velocity  $V$ , the exponents read

$$\begin{aligned} \theta_i &= \epsilon_i \gamma_i (x - Vt + Vt - v_i t - \bar{x}_i) \\ &= \epsilon_i \gamma_i (X_V - (v_i - V)t - \bar{x}_i), \end{aligned} \quad (5.36)$$

where we see the appearance of the “relative velocity”  $v_i - V$ , that is the velocity in the comoving frame.

For each soliton we now have three cases for the limit (5.35), corresponding to a positive, zero or negative relative velocity for the soliton:

Case	$t \rightarrow -\infty$	$t \rightarrow +\infty$
$V < v_i$	$\theta_i \rightarrow +\epsilon_i \infty$	$\theta_i \rightarrow -\epsilon_i \infty$
$V = v_i$	$\theta_i$ finite	$\theta_i$ finite
$V > v_i$	$\theta_i \rightarrow -\epsilon_i \infty$	$\theta_i \rightarrow +\epsilon_i \infty$



Recall that  $\epsilon_i = \pm 1$  is a sign, and  $\gamma_i > 0$  so it does not affect the sign of  $\theta_i$  in the limit.

This tells us that if  $V \neq v_1, v_2$ , then  $\theta_1, \theta_2 \rightarrow \pm\infty$  as  $|t| \rightarrow \infty$ . This implies that<sup>5</sup>

$$\tan \frac{u}{4} = \mu \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}} \rightarrow \pm\infty \text{ or } 0.$$

So  $u/4$  tends to an integer multiple of  $\pi/2$ , which means that  $u$  tends to an integer multiple of  $2\pi$ : the field is in the **vacuum**. The conclusion is that if we go off to infinity in the original  $(x, t)$  plane in any direction apart from  $\frac{dx}{dt} = v_1, v_2$ , then  $u \rightarrow 2\pi n$  for some  $n \in \mathbb{Z}$ .

If instead  $V = v_1$  or  $v_2$ , we need to study the limit more carefully. We will consider a single case  $a_1, a_2 > 0$ , leaving the other cases for the exercises. Since  $a_1 \neq a_2$  for the solution to exist, let us take without loss of generality

$$\boxed{a_2 > a_1 > 0} \implies v_2 > v_1, \quad \epsilon_1 = \epsilon_2 = 1, \quad \mu > 0.$$

Consider  $V = v_1$  first, or "let's ride the slower soliton". In the comoving frame the exponents  $\theta_i$  read

$$\begin{aligned} \theta_1 &= \gamma_1(x - v_1 t - \bar{x}_1) = \gamma_1(X_{v_1} - \bar{x}_1) \\ \theta_2 &= \gamma_2(x - v_2 t - \bar{x}_2) = \gamma_2(X_{v_1} - (v_2 - v_1)t - \bar{x}_2) \end{aligned} \tag{5.37}$$

so  $\theta_1$  stays finite, whereas  $\theta_2 \rightarrow \mp\infty$  as  $t \rightarrow \pm\infty$  with  $X_{v_1}$  fixed (I used that  $v_2 > v_1$ ).

One of the two limits is easier to analyse, so let's start with that:

1.  $t \rightarrow +\infty$ :

In this limit  $\theta_2 \rightarrow -\infty$ , so  $e^{\theta_2} \rightarrow 0$  and

$$\begin{aligned} \tan \frac{u}{4} &= \mu \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}} \\ &\rightarrow \mu e^{\theta_1} \\ &= \mu e^{\gamma_1(X_{v_1} - \bar{x}_1)} \\ &= e^{\gamma_1(x - v_1 t - \bar{x}_1 + \frac{1}{\gamma_1} \log \mu)}, \end{aligned}$$

---

<sup>5</sup>According to the signs of the limits of  $\theta_1$  and  $\theta_2$ , the limit of  $\tan(u/4)$  is as follows:

$$\begin{aligned} ++ : & \quad \tan(u/4) \rightarrow 0 \\ +- : & \quad \tan(u/4) \rightarrow +\infty \\ -+ : & \quad \tan(u/4) \rightarrow -\infty \\ -- : & \quad \tan(u/4) \rightarrow 0. \end{aligned}$$

where in the last line we have expressed the finite limit in the comoving coordinates in terms of the original  $(x, t)$  coordinates.

This is a **kink** (the centre of) which moves with velocity  $v_1$  along the **trajectory**

$$\boxed{x = v_1 t + \bar{x}_1 - \frac{1}{\gamma_1} \log \frac{a_2 + a_1}{a_2 - a_1}}. \quad (5.38)$$

The last term is negative and represents a **backward shift** in space of the slower soliton compared to where it would have been at the same time in the absence of the faster soliton. (Equivalently, we can view this as a time delay for reaching a fixed value of  $x$ .)

2.  $t \rightarrow -\infty$ :

In this limit  $\theta_2 \rightarrow +\infty$ , so  $e^{\theta_2} \rightarrow +\infty$  and

$$\begin{aligned} \tan \frac{u}{4} &= \mu \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}} \\ &\rightarrow -\mu e^{-\theta_1}. \end{aligned}$$

Recalling that  $\tan(A \pm \frac{\pi}{2}) = -\frac{1}{\tan A}$ , this means that

$$\begin{aligned} \tan\left(\frac{u}{4} \pm \frac{\pi}{2}\right) &\rightarrow \mu^{-1} e^{\theta_1} \\ &= e^{\gamma_1 \left(x - v_1 t - \bar{x}_1 - \frac{1}{\gamma_1} \log \mu\right)}. \end{aligned}$$

Therefore

$$u|_{t \rightarrow -\infty, X_{v_1} \text{ finite}} \approx \pm 2\pi + 4 \arctan e^{\gamma_1 \left(x - v_1 t - \bar{x}_1 - \frac{1}{\gamma_1} \log \mu\right)}.$$

(The  $\pm$  sign ambiguity can be fixed by continuity. It turns out that  $-2\pi$  is correct.)

This is a **kink** (the centre of) which moves with velocity  $v_1$  along the **trajectory**


$$\boxed{x = v_1 t + \bar{x}_1 + \frac{1}{\gamma_1} \log \frac{a_2 + a_1}{a_2 - a_1}}. \quad (5.39)$$

The last term is positive and represents a **forward shift** of the slower soliton compared to where it would have been at the same time in the absence of the faster soliton. (Equivalently, we can view this as a time advancement.)

Comparing the trajectories at early times ( $t \rightarrow -\infty$ ) and at late times ( $t \rightarrow +\infty$ ), we see that **the collision with the faster soliton shifts the slower soliton backwards** by

$$\frac{2}{\gamma_1} \log \frac{a_2 + a_1}{a_2 - a_1},$$

as exemplified by this figure:




We say that the slower soliton has a **negative phase shift**:

$$\text{PHASE SHIFT}_{\text{slower}} = -\frac{2}{\gamma_1} \log \frac{a_2 + a_1}{a_2 - a_1} \quad (5.40)$$

We conclude that the slower kink emerges from the collision with the same shape and velocity, but delayed by a finite phase shift.

Now consider  $V = v_2$ , or "let's ride the faster soliton". The calculation is similar to what we did above, so I'll let you work out the details in **[Ex 30]**. If you do this exercise you will find a **surprise**: even though  $a_2 > 0$ , so that acting on the vacuum with the  $a_2$ -Bäcklund transform produces a kink, the component of the two-soliton solution (5.34) that moves at velocity  $v_2$  is actually an **anti-kink**! So, even though the Bäcklund transform always adds a soliton, the nature of the added soliton depends on what is already there.

The shifts have opposite signs to before, as exemplified by this figure:



This results in a **positive phase shift**:

$$\text{PHASE SHIFT}_{\text{faster}} = + \frac{2}{\gamma_2} \log \frac{a_2 + a_1}{a_2 - a_1} . \quad (5.41)$$

Summarising, we have the following picture for the collision of the anti-kink and the kink:




Figure 5.1: Schematic summary of the kink-antikink solution.

See also here for the plot of the kink-antikink solution with parameters  $a_1 = 1.1$  and  $a_2 = 2$ , here for a contour plot of its energy density, which clearly shows the trajectories of the kink and the anti-kink, and here for an animation of the time evolution.

**REMARK:**

From the plot of the exact solution or the contour plot of its energy density we see that **the kink and the anti-kink attract each other**. Indeed we observe that they get closer during the interaction.

The remaining cases for the signs of  $a_1$  and  $a_2$  can be analysed similarly, see [Ex 31] and [Ex 32]. In particular, the 2-soliton solution that contains **two kinks** looks as follows:<sup>6</sup>




Figure 5.2: Schematic summary of the kink-kink solution.

See also here for a plot of the kink-kink solution with parameters  $a_1 = 0.6$  and  $a_2 = -1.5$ , here for a contour plot of its energy density, which clearly shows the trajectories of the two kinks, and here for an animation of the time evolution.

From the plot of the exact solution or the contour plot of its energy density we see that **the two kinks repel each other**. Indeed they get further apart during the interaction. Curiously, they also seem to swap their identities!

**INTERPRETATION:**

ATTRACTIVE FORCE	between	kink and anti-kink
REPULSIVE FORCE	between	kink and kink
REPULSIVE FORCE	between	anti-kink and anti-kink

<sup>6</sup>The solution that contains two anti-kinks can be obtained by sending  $u \mapsto -u$ .

So kinks and anti-kinks behave similarly to elementary particles with electric charge, such as the electron and the positron. The role of electric charge is played here by the topological charge:

Solitons with like topological charges repel  
Solitons with opposite topological charges attract.

It is quite amazing that lump of fields can behave so similarly to pointlike elementary particles. In the 1950's and 1960's, Tony Skyrme used versions of kinks (and anti-kinks) in four spacetime dimensions to model the behaviour of protons and neutrons in atomic nuclei. This is a very far-reaching idea, which unfortunately we don't have time to investigate further in this module.

We have seen that kinks and anti-kinks attract each other. This raises a natural question: can they stick together, or in physics parlance “form a bound state”? The answer is yes. The resulting bound state of a kink and an anti-kink is the “breather”, which we now turn to.

## 5.8 The breather

Recall the general **2-soliton solution** (5.34) of the sine-Gordon equation, that we rewrite here for convenience:

$$u = 4 \arctan \left( \frac{a_2 + a_1}{a_2 - a_1} \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}} \right).$$

This is a solution of the sine-Gordon equation for **any values of** the Bäcklund parameters  $a_1$  **and**  $a_2$  (and integration constants  $c_1$  **and**  $c_2$ ), even **complex** values! However, the sine-Gordon field  $u$  is an angle and so it **must be real**. There are essentially two options to achieve this:<sup>7</sup>

1.  $a_1, a_2$  (and  $c_1, c_2$ )  $\in \mathbb{R}$ : this is what we have considered so far;
2.  $a_2 = \bar{a}_1$  (and  $c_2 = \bar{c}_1$ ): this is what we will consider next. But let's first check that  $u$

---

<sup>7</sup>To be precise, one can also add to the integration constants  $c_1$  and  $c_2$  an integer multiple of  $\pi i$ . This has the effect of permuting the two solitons if the multiple is odd, and has no effect if the multiple is even.

is real:

$$\begin{aligned}
 \bar{u} &= \overline{4 \arctan \left( \frac{a_2 + a_1}{a_2 - a_1} \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}} \right)} \\
 &= 4 \arctan \left( \frac{\bar{a}_2 + \bar{a}_1}{\bar{a}_2 - \bar{a}_1} \frac{e^{\bar{\theta}_1} - e^{\bar{\theta}_2}}{1 + e^{\bar{\theta}_1 + \bar{\theta}_2}} \right) \\
 &= 4 \arctan \left( \frac{a_1 + a_2}{a_1 - a_2} \frac{e^{\theta_2} - e^{\theta_1}}{1 + e^{\theta_2 + \theta_1}} \right) \\
 &= 4 \arctan \left( \frac{a_2 + a_1}{a_2 - a_1} \frac{e^{\theta_1} - e^{\theta_2}}{1 + e^{\theta_1 + \theta_2}} \right) = u .
 \end{aligned}$$

To get to the second line we used the fact that  $\arctan(z)$  and  $e^z$  are complex analytic functions, therefore  $\overline{\arctan(z)} = \arctan(\bar{z})$  and  $\overline{e^z} = e^{\bar{z}}$ . To get to the third line we used  $\theta_2 = \bar{\theta}_1$ , which follows from  $a_2 = \bar{a}_1$  and  $c_2 = \bar{c}_1$ .

Let us then consider option 2 and try a solution with **arbitrary**  $a_1 = \bar{a}_2 \equiv a$  and with  $c_1 = c_2 = 0$  **for simplicity**. Define

$$\begin{aligned}
 a_1 &= a = A + iB = |a|e^{i\varphi} \\
 a_2 &= \bar{a} = A - iB = |a|e^{-i\varphi}
 \end{aligned} \tag{5.42}$$

where  $A = \operatorname{Re}(a)$ ,  $B = \operatorname{Im}(a)$ ,  $\varphi = \arg(a)$ , and let

$$\begin{aligned}
 \theta_1 &= \alpha + i\beta \\
 \theta_2 &= \alpha - i\beta
 \end{aligned} , \tag{5.43}$$

with  $\alpha$  and  $\beta$  real functions of  $x, t$  to be determined below. Then

$$\begin{aligned}
 \tan \frac{u}{4} &= \frac{|a|(e^{-i\varphi} + e^{i\varphi})}{|a|(e^{-i\varphi} - e^{i\varphi})} \cdot \frac{e^{\alpha+i\beta} - e^{\alpha-i\beta}}{1 + e^{2\alpha}} \\
 &= \frac{2 \cos \varphi}{-2i \sin \varphi} \cdot \frac{2i \sin \beta}{2 \cosh \alpha}
 \end{aligned}$$

which simplifies to

$$\tan \frac{u}{4} = -\frac{\cos \varphi}{\sin \varphi} \frac{\sin \beta}{\cosh \alpha} . \tag{5.44}$$

To finish the calculation, let's determine the functions  $\alpha, \beta$  in terms of the coordinates  $x, t$  and the parameters  $|a|$  and  $\varphi$ :

$$\begin{aligned}
 \alpha + i\beta = \theta_1 &= \frac{1}{a}x^+ - ax^- \\
 &= \frac{\bar{a}}{|a|^2}x^+ - ax^- = \frac{A - iB}{|a|^2}x^+ - (A + iB)x^- .
 \end{aligned} \tag{5.45}$$

Therefore

$$\begin{aligned}\alpha = \operatorname{Re}(\theta_1) &= \frac{A}{|a|^2}x^+ - Ax^- \\ &= \frac{A}{|a|} \left( \frac{1}{|a|}x^+ - |a|x^- \right) .\end{aligned}$$

We can now do similar manipulations to what we did after equation (5.15) to find

$$\alpha = \frac{A}{|a|} \gamma(x - vt) \stackrel{(5.42)}{=} \cos \varphi \cdot \gamma(x - vt) , \quad (5.46)$$

where

$$\begin{aligned}v &= \frac{|a|^2 - 1}{|a|^2 + 1} \\ \gamma &= \frac{1}{\sqrt{1 - v^2}} = \frac{1 + |a|^2}{2|a|} .\end{aligned} \quad (5.47)$$

**\* EXERCISE:** Show that similarly **[Ex 33]**

$$\beta = \frac{B}{|a|} \gamma(vx - t) \stackrel{(5.42)}{=} \sin \varphi \cdot \gamma(vx - t) . \quad (5.48)$$

Substituting these expressions in (5.44) we find the **breather** solution

$$\tan \frac{u}{4} = -\cot \varphi \cdot \frac{\sin(\sin \varphi \cdot \gamma(vx - t))}{\cosh(\cos \varphi \cdot \gamma(x - vt))} . \quad (5.49)$$

**REMARK:**

- The ratio of the prefactor and the denominator in the RHS,

$$\frac{-\cot \varphi}{\cosh(\cos \varphi \cdot \gamma(x - vt))} ,$$

defines an **envelope** which moves at the **group velocity**  $v$ . Recall that  $|v| < 1$ , where 1 is the speed of light, so this is consistent with the laws of special relativity.

- The numerator

$$\sin(\sin \varphi \cdot \gamma(x - vt))$$

defines a **carrier wave** which moves at the **phase velocity**  $1/v$ .



To see why the solution (5.49) is called a **“breather”**, let us set  $|a| = 1$ , or equivalently  $v = 0$ . (This can be achieved by switching to a comoving frame if  $v \neq 0$ .) Then the breather simplifies to

$$\tan \frac{u}{4} = \cot \varphi \cdot \frac{\sin(\sin \varphi \cdot t)}{\cosh(\cos \varphi \cdot x)} \quad (5.50)$$

and the field looks like a bouncing (or **“breathing”**) **bound state of a kink and an anti-kink**, with **time period**

$$\tau = \frac{2\pi}{|\sin \varphi|}. \quad (5.51)$$

See figure (5.3) for a summary of the  $v = 0$  breather solution, this for a plot of the breather




Figure 5.3: Schematic summary of the  $v = 0$  breather solution.

solution with  $v = 0$  and  $\varphi = \pi/10$ , this for a contour plot of its energy density, which clearly shows the trajectories of the breathing pair of kink and anti-kink, and this for an animation of the time evolution.

One can show<sup>8</sup> that the  $v = 0$  breather has energy  $E_{\text{breather}} = 16 \cos \varphi$ . Since a static kink and a static anti-kink have energy  $E_{\text{kink}} = E_{\text{antikink}} = 8$ , the binding energy of the kink and the

<sup>8</sup>This is a good but technical exercise, which is not in the problem sheet.

anti-kink in the breather is

$$E_{\text{binding}} = E_{\text{breather}} - E_{\text{kink}} - E_{\text{antikink}} = -16(1 - \cos \varphi) .$$

This is negative as expected: the binding lowers the energy of the solution.

As  $\varphi \rightarrow 0$ , the binding energy tends to zero. It is immediate to see from equation (5.51) that the time period of the bounce diverges:  $\tau \sim 1/|\varphi| \rightarrow \infty$ . The spatial size of the breather also diverges like **[Ex 34]**

$$x_{\text{max}} \sim -\log |\varphi| \rightarrow \infty .$$

In this limit the kink and the antikink become more and more **loosely bound**. The resulting solution

$$u = 4 \arctan (t \cdot \text{sech}(x))$$

describes a kink and an anti-kink starting infinitely far away from one another and doing half an oscillation. Since  $\text{sech}(x) \approx 2e^{-|x|}$  as  $|x| \rightarrow \infty$ , the kink and the anti-kink do not follow linear trajectories as  $t \rightarrow \pm\infty$ . Rather, the asymptotic trajectories of the kink and the anti-kink are given by  $|x| \sim \log |t|$ .

# Chapter 6

## The Hirota method

The main reference for this chapter is §5.3 of [Drazin and Johnson, 1989].

This is an **alternative to the Bäcklund transform** as a way to generate **multi-soliton solutions**, which is sometimes available when the Bäcklund transform is not. It was devised by **Hirota** [Hirota, 1971] to write  **$N$ -soliton solutions of the KdV equation**, and was then generalised to a large class of equations. We will focus on the KdV equation in this chapter.

### 6.1 Motivation

Let us substitute

$$\boxed{u = w_x} \tag{6.1}$$

in the KdV equation

$$u_t + 6uu_x + u_{xxx} = 0 .$$

We find the equation

$$w_{xt} + 6w_x w_{xx} + w_{xxx} = 0 ,$$

which we can integrate with respect to  $x$ :

$$w_t + 3w_x^2 + w_{xxx} = g(t) .$$

We will drop the integration “constant” (with respect to  $x$ )  $g(t)$  in what follows, since it can be absorbed in a redefinition of  $w$  that does not change  $u = w_x$ :

$$w_{\text{old}}(x, t) = w_{\text{new}}(x, t) + \int_{t_0}^t dt' g(t') .$$

Using the new  $w$  (and dropping the subscript “new”), we have the following equation:

$$\boxed{w_t + 3w_x^2 + w_{xxx} = 0} . \quad (6.2)$$

You may ask: why did we make the substitution (6.1)? Recall the **one-soliton solution of KdV**

$$\boxed{u = 2\mu^2 \operatorname{sech}^2 [\mu(x - x_0 - 4\mu^2 t)]} \quad (6.3)$$

with

$$\boxed{\mu = \frac{\sqrt{v}}{2}} . \quad (6.4)$$

This one-soliton solution can be written as  $u = w_x$  with

$$\boxed{w = 2\mu \tanh [\mu(x - x_0 - 4\mu^2 t)]} . \quad (6.5)$$

In fact, we can integrate the right-hand side of (6.5) once more, using

$$\tanh y = \frac{d}{dy} \log \cosh y .$$

Therefore the one-soliton solution (6.3) of KdV can be written as

$$\boxed{u = 2 \frac{\partial^2}{\partial x^2} \log \cosh [\mu(x - x_0 - 4\mu^2 t)]} . \quad (6.6)$$

This can be simplified further. Let

$$\boxed{X = x - x_0 - 4\mu^2 t} . \quad (6.7)$$

Then

$$\begin{aligned} u &= 2 \frac{\partial^2}{\partial X^2} \log \frac{e^{-\mu X} (1 + e^{2\mu X})}{2} \\ &= 2 \frac{\partial^2}{\partial X^2} [-\mu X - \log 2 + \log (1 + e^{2\mu X})] \\ &= 2 \frac{\partial^2}{\partial X^2} \log (1 + e^{2\mu X}) , \end{aligned}$$

or in terms of the original coordinates

$$\boxed{u(x, t) = 2 \frac{\partial^2}{\partial x^2} \log (1 + e^{2\mu(x - x_0 - 4\mu^2 t)})} . \quad (6.8)$$

This is the form of the **one-soliton solution of KdV** that we will refer to in the following.

## 6.2 KdV equation in bilinear form

### 6.2.1 The quadratic form of KdV

Inspired by the form (6.8) of the one-soliton solution, let's substitute

$$\boxed{w = 2 \frac{\partial}{\partial x} \log f = \frac{f_x}{f}} \iff \boxed{u = 2 \frac{\partial^2}{\partial x^2} \log f} \quad (6.9)$$

in equation (6.2).<sup>1</sup> Then

$$\begin{aligned} \frac{1}{2}w_t &= \frac{f_{xt}f - f_x f_t}{f^2}, \\ \frac{1}{2}w_x &= \frac{f_{xx}f - f_x^2}{f^2}, \\ \frac{1}{2}w_{xx} &= \dots \quad \text{[Ex 35]} \\ \frac{1}{2}w_{xxx} &= \frac{f_{xxx}f}{f} - 4 \frac{f_{xxx}f_x}{f^2} - 3 \frac{f_{xx}^2}{f^2} + 12 \frac{f_{xx}f_x^2}{f^3} - 6 \frac{f_x^4}{f^4}, \end{aligned} \quad (6.10)$$

and equation (6.2) for  $w$  becomes [Ex 35]

$$\frac{f_{xt}}{f} - \frac{f_x f_t}{f^2} + 3 \frac{f_{xx}^2}{f^2} - 4 \frac{f_{xxx}f_x}{f^2} + \frac{f_{xxxx}}{f} = 0$$

for  $f$ .

Multiplying by  $f^2$ , we find the so called **quadratic form of the KdV equation**:

$$\boxed{f f_{xt} - f_x f_t + 3f_{xx}^2 - 4f_x f_{xxx} + f f_{xxxx} = 0}. \quad (6.11)$$

Some cancellations have taken place to get to the quadratic form (6.11) of the KdV equation, but at first sight this might not seem progress on the initial equation (6.2). But (6.11) is **quadratic in  $f$**  and it **can be rewritten in a neat way**. A hint for that is that

$$\frac{\partial}{\partial x} \frac{\partial}{\partial t} \left( \frac{1}{2} f^2 \right) = \frac{\partial}{\partial x} (f f_t) = f f_{xt} + f_x f_t.$$

This is almost like the first two terms in (6.11), except for the relative sign. We will fix this sign problem shortly.

---

<sup>1</sup>In the literature on integrable systems, the function  $f$  is now called the  $\tau$ -function.

### 6.2.2 Hirota's bilinear operator

Hirota defined a **bilinear differential operator**  $D$  which maps a **pair** of functions  $(f, g)$  into a **single** function  $D(f, g)$ . If we work on  $C^\infty$  functions, then

$$D : \quad C^\infty \times C^\infty \rightarrow C^\infty$$

$$(f, g) \mapsto D(f, g) ,$$

and bilinearity means that

$$D(a_1 f_1 + a_2 f_2, g) = a_1 D(f_1, g) + a_2 D(f_2, g)$$

$$D(f, b_1 g_1 + b_2 g_2) = b_1 D(f, g_1) + b_2 D(f, g_2)$$

for any constants  $a_1, a_2, b_1, b_2$ .

**REMARK:**

This is unlike the usual linear differential operators that you are familiar with, such as  $\left(\frac{\partial}{\partial x}\right)^n$ , which maps a single function  $f$  into a single function  $\frac{\partial^n f}{\partial x^n}$ .

For any integers  $m, n \geq 0$ , we **define** the bilinear differential operators  $D_t^m D_x^n$  by

$$\boxed{[D_t^m D_x^n(f, g)](x, t) := \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'}\right)^m \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'}\right)^n f(x, t)g(x', t') \Big|_{\substack{x'=x \\ t'=t}}} .} \quad (6.12)$$

Let us look at a few examples. We start with

$$\begin{aligned} [D_t(f, g)](x, t) &= \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'}\right) f(x, t)g(x', t') \Big|_{\substack{x'=x \\ t'=t}} \\ &= f_x(x, t)g(x', t') - f(x, t)g_{t'}(x', t') \Big|_{\substack{x'=x \\ t'=t}} \\ &= f_t(x, t)g(x, t) - f(x, t)g_t(x, t) , \end{aligned} \quad (6.13)$$

so

$$D_t(f, g) = f_t g - f g_t \quad \text{and} \quad D_t(f, f) = 0 ,$$

and similarly for  $D_x$ . Next we look at

$$\begin{aligned} [D_t D_x(f, g)](x, t) &= \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'}\right) \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'}\right) f(x, t)g(x', t') \Big|_{\substack{x'=x \\ t'=t}} \\ &= \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'}\right) (f_x(x, t)g(x', t') - f(x, t)g_{x'}(x', t')) \Big|_{\substack{x'=x \\ t'=t}} \\ &= f_{xt}(x, t)g(x, t) - f_t(x, t)g_x(x, t) - f_x(x, t)g_t(x, t) + f(x, t)g_{xt}(x, t) , \end{aligned} \quad (6.14)$$

so

$$D_t D_x(f, g) = f_{xt}g - f_t g_x - f_x g_t + f g_{xt} \quad \text{and} \quad D_t D_x(f, f) = 2(f f_{tx} - f_t f) . \quad (6.15)$$

This is promising, because the right-hand-side of the last expression reproduces the first two terms in the quadratic form of the KdV equation (6.11), up to an overall factor of 2. Let's proceed then and compute

$$D_x^2(f, g) = f_{xx}g - 2f_x g_x + f g_{xx} , \quad (6.16)$$

which implies

$$D_x^2(f, f) = 2(f f_{xx} - f_x^2) .$$

**REMARK:**

Note that  $D_x^2(f, f) \neq 0$  even though  $D_x(f, f) = 0$ . This is not inconsistent, because  $D_x^2(f, f) \neq D_x(D_x(f, f))$ . In fact, the right-hand side of this last expression is meaningless, since the outer  $D_x$  must act on a pair of functions, but  $D_x(f, f)$  is a single function.

Finally, we can calculate

$$\begin{aligned} D_x^4(f, g) &= \dots & \text{[Ex 36]} \\ &= f_{xxxx}g - 4f_{xxx}g_x + 6f_{xx}g_{xx} - 4f_x g_{xxx} + f g_{xxxx} . \end{aligned} \quad (6.17)$$

Note that the result is like  $\partial_x^4(fg)$ , but with alternating signs! So

$$D_x^4(f, f) = 2(f f_{xxxx} - 4f_x f_{xxx} + 3f_{xx}^2) . \quad (6.18)$$

Here is the **miracle**: the KdV equation in its quadratic form (6.11) can be recast as

$$\boxed{(D_t D_x + D_x^4)(f, f) = 0} \quad (6.19)$$

where the bilinear operator  $D_t D_x + D_x^4$  is defined by linearity on the space of operators of the type (6.12), namely  $(D_t D_x + D_x^4)(f, g) = D_t D_x(f, g) + D_x^4(f, g)$ . Equation (6.19) is the so called **bilinear form of the KdV equation**.

**REMARK:**

Observe that we can formally factor the Hirota operator as

$$D_t D_x + D_x^4 = (D_t + D_x^3)D_x ,$$

which is a short-hand for

$$(D_t D_x + D_x^4)(f, g) = (\partial_t - \partial_{t'} + (\partial_x - \partial_{x'})^3)(\partial_x - \partial_{x'})f(x, t)g(x', t') \Big|_{\substack{x'=x \\ t'=t}} .$$

This is not an accident. It is related to the fact that the differential operator  $\partial_t + \partial_x^3$  appears in the linearised KdV equation for  $u$ , and therefore the differential operator  $(\partial_t + \partial_x^3)\partial_x$  appears in the linearisation of the equation for  $w$  (before integration with respect to  $x$ ).

## 6.3 Solutions

We will need **two ideas** to find multi-soliton solutions. The **first idea** is inspired by a rather basic observation: if we take  $f = 1$ , then the KdV field is the vacuum  $u = 0$ ; if instead we take

$$f = 1 + e^{2\mu(x-x_0-4\mu^2t)},$$

then the KdV field  $u$  is the one-soliton solution (6.8). Since (6.19) is a **bilinear** equation, this suggests that **multi-soliton solutions** might be obtained from an  $f$  which is a **sum of exponentials of linear functions of  $x$  and  $t$** , with  $1 = e^0$  as the trivial case. But before we get to the general case, let us check the Hirota formalism by rederiving the one-soliton solution of the KdV equation.

### 6.3.1 Example: 1-soliton

Let's try

$$\boxed{f = 1 + e^\theta} \quad (6.20)$$

with

$$\boxed{\theta = ax + bt + c}, \quad (6.21)$$

where  $a, b, c$  are constants.

**Lemma 1.** *If  $\theta_i = a_i x + b_i t + c_i$  ( $i = 1, 2$ ), then [Ex 38]*

$$\boxed{D_t^m D_x^n (e^{\theta_1}, e^{\theta_2}) = (b_1 - b_2)^m (a_1 - a_2)^n e^{\theta_1 + \theta_2}}. \quad (6.22)$$

In particular

$$\begin{aligned} D_t^m D_x^n (e^\theta, e^\theta) &= 0 && \text{(unless } m = n = 0) \\ D_t^m D_x^n (e^\theta, 1) &= (-1)^{m+n} D_t^m D_x^n (1, e^\theta) = b^m a^n e^\theta. \end{aligned} \quad (6.23)$$

Therefore the bilinear form of the KdV equation for  $f = 1 + e^\theta$  is

$$\begin{aligned} 0 &= (D_t D_x + D_x^4)(1 + e^\theta, 1 + e^\theta) \\ &\stackrel{\text{bilinearity}}{=} (D_t D_x + D_x^4) [(1, 1) + (1, e^\theta) + (e^\theta, 1) + (e^\theta, e^\theta)] \\ &\stackrel{(6.23)}{=} 2(D_t D_x + D_x^4)(e^\theta, 1) \\ &\stackrel{(6.23)}{=} 2(ba + a^4)e^\theta = 2a(b + a^3)e^\theta. \end{aligned} \quad (6.24)$$

There are two ways to solve this algebraic equation:



1.  $\underline{a = 0}$ : then  $f$  is independent of  $x$ , and  $u = 0$ .

2.  $\underline{b = -a^3}$ : then

$$f = 1 + e^{ax - a^3t + c},$$

and

$$u = 2 \frac{\partial^2}{\partial x^2} \log \left( 1 + e^{ax - a^3t + c} \right), \quad (6.25)$$

which is nothing but the one-soliton solution (6.8), up to a redefinitions of the constants.

### 6.3.2 $N$ -soliton solutions

The **second idea** is to look for a **power series solution** (or a so called “perturbative expansion”) in an **auxiliary parameter**  $\epsilon$ ,

$$f(x, t) = \sum_{n=0}^{\infty} \epsilon^n f_n(x, t) \quad \text{with} \quad f_0 = 1, \quad (6.26)$$

and hope that the series terminates at some value of  $n$ , so that we can take  $\epsilon$  to be finite and eventually set it to 1.

We will write the bilinear form of KdV as

$$B(f, f) = 0 \quad \text{with} \quad B = D_t D_x + D_x^4. \quad (6.27)$$

Substituting (6.26) in (6.27), we find

$$\begin{aligned} 0 &= B\left(\sum_{n_1=0}^{\infty} \epsilon^{n_1} f_{n_1}, \sum_{n_2=0}^{\infty} \epsilon^{n_2} f_{n_2}\right) \\ &= \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \epsilon^{n_1+n_2} B(f_{n_1}, f_{n_2}) \end{aligned} \quad (6.28)$$

where in the second line we used the bilinearity of the Hirota operator  $B$ . Gathering terms of the same degree  $n = n_1 + n_2$  in  $\epsilon$ , we can rewrite (6.28) as

$$0 = \sum_{n=0}^{\infty} \epsilon^n \sum_{m=0}^n B(f_{n-m}, f_m) \stackrel{B(1,1)=0}{=} \sum_{n=1}^{\infty} \epsilon^n \sum_{m=0}^n B(f_{n-m}, f_m). \quad (6.29)$$

Let's solve this equation order by order in  $\epsilon$ . We find that

$$\sum_{m=0}^n B(f_{n-m}, f_m) = 0 \quad \forall n = 1, 2, \dots \quad (6.30)$$

with  $f_0 = 1$ . Writing (6.30) as

$$\boxed{B(f_n, 1) + B(1, f_n) = (\text{expression in } f_1, f_2, \dots, f_{n-1})}, \quad (6.31)$$

makes it clear that we can **solve** (6.30) **recursively** to determine the Taylor coefficients of  $f$ . We will need another lemma:

**Lemma 2.** [Ex 39] *For any function  $f$ ,*

$$\boxed{D_t^m D_x^n (f, 1) = (-1)^{m+n} D_t^m D_x^n (1, f) = \frac{\partial^m}{\partial t^m} \frac{\partial^n}{\partial x^n} f}. \quad (6.32)$$

Using this lemma, we can write the recursion relation (6.31) more explicitly as

$$\boxed{\frac{\partial}{\partial x} \left( \frac{\partial}{\partial t} + \frac{\partial^3}{\partial x^3} \right) f_n = -\frac{1}{2} \sum_{m=1}^{n-1} B(f_{n-m}, f_m)}, \quad (6.33)$$

which is valid for all  $n = 1, 2, \dots$ . In the following I will refer to this recursion relation, which determines  $f_n$  in terms of all the  $f_m$  with  $m < n$ , as Eq <sub>$n$</sub> .

For  $n = 1$  this reduces to

$$\frac{\partial}{\partial x} \left( \frac{\partial}{\partial t} + \frac{\partial^3}{\partial x^3} \right) f_1 = 0$$

or, with appropriate boundary conditions,

$$\boxed{\left( \frac{\partial}{\partial t} + \frac{\partial^3}{\partial x^3} \right) f_1 = 0}, \quad (6.34)$$

which is a linear equation. A simple solution is

$$\boxed{f_1 = \sum_{i=1}^N e^{a_i x - a_i^3 t + c_i} \equiv \sum_{i=1}^N e^{\theta_i}}, \quad (6.35)$$

where  $a_i$  and  $c_i$  are constants as usual.

The higher  $f_n$  are then determined recursively using Eq <sub>$n$</sub>  (6.33). The **amazing fact** is that with  $f_1$  as in equation (6.35), **the expansion (6.26) terminates at order  $N$** . All the higher equations Eq <sub>$n > N$</sub>  are solved with  $f_{n > N} = 0$ ! This is quite non-trivial: it requires that  $f_1, \dots, f_N$  satisfy the consistency conditions that the RHS of Eq <sub>$n$</sub>  vanish for  $n = N + 1, \dots, 2N$ .

We then find that the  **$N$ -soliton solution of KdV** is given by

$$\boxed{f = 1 + f_1 + f_2 + \dots + f_N}, \quad (6.36)$$

where we set  $\epsilon = 1$  (or absorbed it in the constants  $c_i$ ).

### EXAMPLES:

$N = 1$  In this case

$$f_1 = e^{a_1 x - a_1^3 t + c_1} \equiv e^{\theta_1}$$

and Eq<sub>2</sub> reads

$$\partial_x(\partial_t + \partial_x^3)f_2 = -\frac{1}{2}B(e^{\theta_1}, e^{\theta_1}) \stackrel{(6.23)}{=} 0.$$

So we can take  $f_2 = 0$  (and  $f_3 = f_4 = \dots = 0$  as well). Setting  $\epsilon = 1$ , or absorbing  $\epsilon$  in  $c_1$ , we get

$$f = 1 + e^{\theta_1},$$

the one-soliton solution as we found in (6.25).

$N = 2$  In this case

$$f_1 = e^{\theta_1} + e^{\theta_2}$$

and Eq<sub>2</sub> reads

$$\begin{aligned} \partial_x(\partial_t + \partial_x^3)f_2 &= -\frac{1}{2}B(e^{\theta_1} + e^{\theta_2}, e^{\theta_1} + e^{\theta_2}) \\ &\stackrel{\text{bilinearity}}{=} -B(e^{\theta_1}, e^{\theta_2}) \\ &\stackrel{B=D_t D_x + D_x^4}{+ (6.22)} -(a_1 - a_2)[-a_1^3 + a_2^3 + (a_1 - a_2)^3]e^{\theta_1 + \theta_2} \\ &\stackrel{+ (6.22)}{=} 3a_1 a_2 (a_1 - a_2)^2 e^{\theta_1 + \theta_2}. \end{aligned} \tag{6.37}$$

So let's try

$$f_2 = A e^{\theta_1 + \theta_2}$$

for some constant  $A$  to be determined. Substituting in the previous equation we find

$$\begin{aligned} (a_1 + a_2)[-a_1^3 - a_2^3 + (a_1 + a_2)^3]A e^{\theta_1 + \theta_2} &= 3a_1 a_2 (a_1 - a_2)^2 e^{\theta_1 + \theta_2} \\ 3a_1 a_2 (a_1 + a_2)^2 A &= 3a_1 a_2 (a_1 - a_2)^2 \\ A &= \left( \frac{a_1 - a_2}{a_1 + a_2} \right)^2. \end{aligned} \tag{6.38}$$

So we get

$$f = 1 + e^{\theta_1} + e^{\theta_2} + \left( \frac{a_1 - a_2}{a_1 + a_2} \right)^2 e^{\theta_1 + \theta_2} \tag{6.39}$$

for the **2-soliton solution of KdV**, where again we set  $\epsilon = 1$  or absorbed into a shift of  $c_1$  and  $c_2$ .

\* **EXERCISE:** Show that  $B(f_1, f_2) = 0$  and  $B(f_2, f_2) = 0$ , so that one can consistently set  $f_3 = f_4 = \dots = 0$ . **[Ex 40]**

General  $N$  Let's first massage the 2-soliton solution (6.39) that we have just found:

$$\begin{aligned} f &= (1 + e^{\theta_1})(1 + e^{\theta_2}) - e^{\theta_1 + \theta_2} + \left( \frac{a_1 - a_2}{a_1 + a_2} \right)^2 e^{\theta_1 + \theta_2} \\ &= (1 + e^{\theta_1})(1 + e^{\theta_2}) - \frac{4a_1 a_2}{(a_1 + a_2)^2} e^{\theta_1 + \theta_2} \\ &= \begin{vmatrix} 1 + e^{\theta_1} & \frac{2a_1}{a_1 + a_2} e^{\theta_2} \\ \frac{2a_2}{a_1 + a_2} e^{\theta_1} & 1 + e^{\theta_2} \end{vmatrix}. \end{aligned}$$

So we can write

$$\boxed{f = \det(S)}, \quad \text{where} \quad \boxed{S_{ij} = \delta_{ij} + \frac{2a_i}{a_i + a_j} e^{\theta_j}}, \quad (6.40)$$

where here  $i, j \in \{1, 2\}$ .<sup>2</sup>

It turns out that formula (6.40) generalises to higher  $N$ , with  $S$  an  $N \times N$  matrix of the same form as in (6.40) but with  $i, j \in \{1, \dots, N\}$ , giving the  $N$ -**soliton solution of KdV**. This can be proven by induction. One can also show that

$$f_n = \sum_{1 \leq i_1 < i_2 < \dots < i_n \leq N} e^{\theta_{i_1} + \theta_{i_2} + \dots + \theta_{i_n}} \prod_{1 \leq j < k \leq n} \left( \frac{a_{i_j} - a_{i_k}}{a_{i_j} + a_{i_k}} \right)^2.$$

## 6.4 Asymptotics of 2-soliton solutions and phase shifts

To see that the  $N = 2$  solution (6.39) does indeed involve two solitons, we can follow the same logic of section 5.7, where we studied the asymptotics of 2-soliton solutions of the sine-Gordon equation. Namely, we switch to an appropriate comoving frame and only then take  $t \rightarrow \pm\infty$ .

Recall that

$$f = 1 + e^{\theta_1} + e^{\theta_2} + A e^{\theta_1 + \theta_2}$$

where

$$\theta_i = a_i x - a_i^3 t + c_i, \quad A = \left( \frac{a_1 - a_2}{a_1 + a_2} \right)^2.$$

---

<sup>2</sup>Note that using  $e^{\theta_i}$  instead of  $e^{\theta_j}$  in the definition of the matrix element  $S_{ij}$  produces the same determinant.

We can take  $0 < a_1 < a_2$  without loss of generality<sup>3</sup> so  $v_1 = a_1^2 < v_2 = a_2^2$ . Let's **follow the slower soliton** first:

$$\boxed{t \rightarrow \pm\infty \quad \text{with} \quad X_{a_1^2} = x - a_1^2 t \quad \text{fixed}} . \quad (6.41)$$

Then

$$\begin{aligned} \theta_1 &= a_1 X_{a_1^2} + c_1 \\ \theta_2 &= a_2 \left( X_{a_1^2} - (a_2^2 - a_1^2)t \right) + c_2 . \end{aligned} \quad (6.42)$$

Let us consider the two limits (6.41) in turn.

1.  $t \rightarrow +\infty$ : in this limit  $\theta_1$  stays fixed and  $\theta_2 \rightarrow -\infty$ , so

$$f \equiv 1 + e^{\theta_1} . \quad (6.43)$$

This describes a KdV soliton centred at

$$\boxed{x_{\text{centre}}(t) = a_1^2 t - \frac{c_1}{a_1}} . \quad (6.44)$$

2.  $t \rightarrow -\infty$ : in this limit  $\theta_1$  stays fixed and  $\theta_2 \rightarrow +\infty$ , so

$$f \equiv e^{\theta_2} (1 + A e^{\theta_1}) . \quad (6.45)$$

The prefactor  $e^{\theta_2}$  does not matter, because

$$\begin{aligned} u &= 2 \frac{\partial^2}{\partial x^2} \log f \equiv 2 \frac{\partial^2}{\partial x^2} [\theta_2 + \log(1 + A e^{\theta_1})] \\ &= 2 \frac{\partial^2}{\partial x^2} \log(1 + A e^{\theta_1}) \\ &= 2 \frac{\partial^2}{\partial x^2} \log \left( 1 + e^{a_1 x - a_1^3 t + c_1 + \log A} \right) . \end{aligned} \quad (6.46)$$

where in the second line we used that  $\theta_2$  is linear in  $x$ , and in the third line we expressed the result in the original  $(x, t)$  coordinates. This describes a KdV soliton centred at

$$\boxed{x_{\text{centre}}(t) = a_1^2 t - \frac{c_1 + \log A}{a_1}} . \quad (6.47)$$

Therefore the **slower soliton** has a **negative phase shift**:

$$\boxed{\text{PHASE SHIFT}_{\text{slower}} = \frac{1}{a_1} \log A = -\frac{2}{a_1} \log \left| \frac{a_2 + a_1}{a_2 - a_1} \right| < 0} . \quad (6.48)$$

---

<sup>3</sup>Convince yourself of this statement.

Next, let's **follow the faster soliton**:

$$\boxed{t \rightarrow \pm\infty \quad \text{with} \quad X_{a_2^2} = x - a_2^2 t \quad \text{fixed}} . \quad (6.49)$$

Then

$$\begin{aligned} \theta_1 &= a_1 \left( X_{a_2^2} - (a_1^2 - a_2^2)t \right) + c_1 \\ \theta_2 &= a_2 X_{a_2^2} + c_2 . \end{aligned} \quad (6.50)$$

Let us consider the two limits (6.49) in turn.

1.  $t \rightarrow -\infty$ : in this limit  $\theta_1 \rightarrow -\infty$  and  $\theta_2$  stays fixed, so

$$f \equiv 1 + e^{\theta_2} .$$

This describes a KdV soliton centred at

$$\boxed{x_{\text{centre}}(t) = a_2^2 t - \frac{c_2}{a_2}} . \quad (6.51)$$

2.  $t \rightarrow +\infty$ : in this limit  $\theta_1 \rightarrow +\infty$  and  $\theta_2$  stays fixed, so

$$f \equiv e^{\theta_1} (1 + A e^{\theta_2}) ,$$

which describes a KdV soliton centred at

$$\boxed{x_{\text{centre}}(t) = a_2^2 t - \frac{c_2 + \log A}{a_2}} . \quad (6.52)$$

Therefore the **faster soliton** has a **positive phase shift**:

$$\boxed{\text{PHASE SHIFT}_{\text{faster}} = -\frac{1}{a_2} \log A = \frac{2}{a_2} \log \left| \frac{a_2 + a_1}{a_2 - a_1} \right| > 0} . \quad (6.53)$$

Summarising, from the analysis of the asymptotics of the 2-soliton solution we obtain the picture in Fig. 6.1. We have therefore verified that KdV solitons satisfy the third defining property of a soliton 3: when two KdV solitons collide, they emerge from the collision with the same shapes and velocities that they had before the collision. The effect of the interaction is in the phase shifts of the two solitons, which capture the advancement of the faster soliton and the delay of the slower soliton.

We can also look at the exact 2-soliton solution (6.9) and (6.39) to get a better feel for what happens during the collision. Here is a plot of the 2-soliton solution with parameters  $a_1 = 0.7$  and  $a_2 = 1$ . The contour plot of its energy density clearly shows the trajectories of the two KdV solitons and how they repel each other and swap identities when they get close. Finally, here is an animation of their time evolution.




Figure 6.1: Schematic summary of the 2-soliton solution of KdV.

# Bibliography

Boussinesq, J. (1877). *Essai sur la théorie des eaux courantes*. Impr. nationale.

Dauxois, T. (2008). Fermi, Pasta, Ulam and a mysterious lady. *Physics Today*, 61(1):55–57.  
arXiv: 0801.1590.

Dauxois, T. and Peyrard, M. (2006). *Physics of Solitons*. Cambridge University Press.

Drazin, P. G. and Johnson, R. S. (1989). *Solitons: An Introduction*. Cambridge University Press.

Fermi, E., Pasta, J. R., Ulam, S. M., and Tsingou, M. (1955). Studies of nonlinear problems. No. LA-1940. Los Alamos Scientific Lab., N. Mex.

Gordon, W. (1926). Der Comptoneffekt nach der Schrödingerschen Theorie. *Zeitschrift für Physik*, 40(1):117–133.

Hirota, R. (1971). Exact solution of the Korteweg—de Vries equation for multiple collisions of solitons. *Phys. Rev. Lett.*, 27:1192–1194.

John Scott Russell (1845). *Report on Waves: Made to the Meetings of the British Association in 1842-43*.

Klein, O. (1926). Quantentheorie und fünfdimensionale Relativitätstheorie. *Zeitschrift für Physik*, 37(12):895–906.

Korteweg, D. D. J. and Vries, D. G. d. (1895). On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 39(240):422–443.



- Manton, N. and Sutcliffe, P. (2004). *Topological Solitons*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge.
- Miura, R. M. (1968). Korteweg-de Vries Equation and Generalizations. I. A Remarkable Explicit Nonlinear Transformation. *Journal of Mathematical Physics*, 9(8):1202–1204.
- Miura, R. M., Gardner, C. S., and Kruskal, M. D. (1968). Korteweg – de Vries Equation and Generalizations. II. Existence of Conservation Laws and Constants of Motion. *Journal of Mathematical Physics*, 9(8):1204–1209.
- Porter, M. A., Zabusky, N. J., Hu, B., and Campbell, D. K. (2009). Fermi, Pasta, Ulam and the birth of experimental mathematics: a numerical experiment that Enrico Fermi, John Pasta, and Stanislaw Ulam reported 54 years ago continues to inspire discovery. *American Scientist*, 97(3):214–221.
- Takahashi, D. and Satsuma, J. (1990). A soliton cellular automaton. 59:3514–3519.
- Zabusky, N. J. and Kruskal, M. D. (1965). Interaction of “solitons” in a collisionless plasma and the recurrence of initial states. *Physical review letters*, 15(6):240.